

# TIKHONOV REGULARIZATION AND TOTAL LEAST SQUARES

GENE H. GOLUB\*, PER CHRISTIAN HANSEN† AND DIANNE P. O'LEARY‡

**Abstract.** Discretizations of inverse problems lead to systems of linear equations with a highly ill-conditioned coefficient matrix, and in order to compute stable solutions to these systems it is necessary to apply regularization methods. We show how Tikhonov's regularization method, which in its original formulation involves a least squares problem, can be recast in a total least squares formulation, suited for problems in which both the coefficient matrix and the right-hand side are known only approximately. We analyze the regularizing properties of this method and demonstrate by a numerical example that in certain cases with large perturbations, the new method is superior to standard regularization methods.

**Key words.** total least squares, discrete ill-posed problems, regularization, bidiagonalization.

**AMS(MOS) subject classifications.** 65F20, 65F30.

**1. Introduction.** In this paper we study a class of methods for producing an approximate solution to a linear system of equations  $Ax \approx b$  where  $A$  is  $m \times n$  with  $m > n$ . We assume that the elements of  $A$  and those of  $b$  are contaminated by some noise. An appropriate statement of the problem in the case of noisy data is the *total least squares* (TLS) formulation [5], [6, §12.3], [16]:

$$(1) \quad \min_{\tilde{A}, \tilde{b}, x} \|(A, b) - (\tilde{A}, \tilde{b})\|_F \quad \text{subject to} \quad \tilde{b} = \tilde{A}x.$$

In contrast, if the elements of  $A$  are exact and only  $b$  contains noise, then the corresponding formulation is the *least squares* (LS) problem

$$(2) \quad \min_x \|Ax - b\|_2$$

Our focus in this work is on very ill-conditioned problems where the singular values of  $A$  decay gradually to zero. Such problems arise, for example, from the discretization of ill-posed problems such as integral equations of the first kind. See, e.g., [2], [3], [7], [8], and [10] for examples and details.

In these problems, the solutions to the two formulations in (1) and (2) can be hopelessly contaminated by the noise in directions corresponding to small singular values of  $A$  or  $(A, b)$ . Because of this, it is necessary to compute a *regularized solution* in which the effect of such noise is filtered out. For least squares problems, the filtering is often done by Tikhonov's regularization method where a damping is added to each SVD component of the solution, thus effectively filtering out the components corresponding to the small singular values. Alternatively, one can truncate the SVD expansion of the solution, leaving out all the SVD components corresponding to the small singular values.

For the total least squares problem, the truncation approach has already been studied by Fierro, Golub, Hansen, and O'Leary [4]. In the present work we focus on

---

\* Department of Computer Science, Stanford University, Stanford, CA 94305 (golub@sccm.stanford.edu).

† Department of Mathematical Modelling, Technical University of Denmark, Building 321, DK-2800 Lyngby, Denmark (pch@imm.dtu.dk).

‡ Department of Computer Science and Institute for Advanced Computer Studies, University of Maryland, College Park, MD 20742 (oleary@cs.umd.edu). The work of this author was supported by the National Science Foundation under Grant CCR 95-03126 and by a General Research Board Grant from the Office of Graduate Studies and Research of the University of Maryland.

the Tikhonov regularization approach for total least squares. We thus arrive at a new regularization method in which stabilization enters the formulation in a natural way, and that is able to produce regularized solutions with superior properties for certain problems in which the perturbations are large.

Our paper is organized as follows. In §2 we introduce the regularized TLS method, and we study its regularizing properties. Computational aspects are described in §3, and we conclude the paper with a numerical example in §4. A preliminary report on this work appeared as [12].

**2. The Regularized TLS Method.** Our regularization of the TLS problem is based on Tikhonov regularization. For the linear least squares problem (2), a general version of Tikhonov’s method takes the form

$$(3) \quad \min \{ \|Ax - b\|_2^2 + \lambda \|Lx\|_2^2 \} ,$$

where  $\lambda$  is a positive constant chosen to control the size of the solution vector, and  $L$  is a matrix that defines a (semi)norm on the solution through which the “size” is measured [11]. Often,  $L$  represents the first or second derivative operator. If  $L$  is the identity matrix, then the Tikhonov problem is said to be in standard form. The solution  $x_\lambda$  to (3) solves the problem

$$(4) \quad (A^T A + \lambda L^T L)x = A^T b.$$

As  $\lambda$  increases, the (semi)norm  $\|Lx\|_2$  of the solution vector decreases monotonically while the residual  $\|Ax - b\|_2$  increases monotonically.

Tikhonov regularization has an important equivalent formulation as

$$(5) \quad \min \|Ax - b\|_2 \quad \text{subject to} \quad \|Lx\|_2 \leq \delta ,$$

where  $\delta$  is a positive constant. Problem (5) is a least squares problem with a quadratic constraint, and using the Lagrange multiplier formulation

$$(6) \quad \mathcal{L}(x, \lambda) = \|Ax - b\|_2^2 + \lambda (\|Lx\|_2^2 - \delta^2) ,$$

it can be shown that if  $\delta \leq \|x_{\text{LS}}\|_2$ , where  $x_{\text{LS}}$  is the least squares solution to (2), then the solution  $x_\delta$  to (5) is identical to the solution  $x_\lambda$  to (3) for an appropriately chosen  $\lambda$ , and there is a monotonic relation between the parameters  $\delta$  and  $\lambda$ .

To carry this idea over to the TLS setting, we add an upper bound on the (semi)norm  $\|Lx\|_2$  of the solution vector  $x$  in the TLS problem (1). The formulation of the regularized TLS (R-TLS) problem thus becomes

$$(7) \quad \min \|(A, b) - (\tilde{A}, \tilde{b})\|_F \quad \text{subject to} \quad \tilde{b} = \tilde{A}x , \quad \|Lx\|_2 \leq \delta ,$$

and the corresponding Lagrange multiplier formulation is

$$(8) \quad \hat{\mathcal{L}}(\tilde{A}, x, \mu) = \|(A, b) - (\tilde{A}, \tilde{A}x)\|_F^2 + \mu (\|Lx\|_2^2 - \delta^2) ,$$

where  $\mu$  is the Lagrange multiplier, zero if the inequality constraint is inactive. The solution  $\tilde{x}_\delta$  to this problem is different from the solution  $x_{\text{TLS}}$  to (1) whenever  $\delta$  is less than  $\|Lx_{\text{TLS}}\|_2$ . The two solutions  $\tilde{x}_\delta$  and  $x_\delta$  to the two regularized problems in (5) and (7) have a surprising relationship, explained by the following theorem.

**THEOREM 2.1.** *The regularized TLS solution  $\tilde{x}_\delta$  to (7), with the inequality constraint replaced by equality, is a solution to the problem*

$$(9) \quad (A^T A + \lambda_I I_n + \lambda_L L^T L)x = A^T b$$

where the parameters  $\lambda_I$  and  $\lambda_L$  are given by

$$(10) \quad \lambda_I = -\frac{\|b - Ax\|_2^2}{1 + \|x\|_2^2}$$

$$(11) \quad \lambda_L = \mu (1 + \|x\|_2^2)$$

and where  $\mu$  is the Lagrange multiplier in (8). The two parameters are related by

$$(12) \quad \lambda_L \delta^2 = b^T (b - Ax) + \lambda_I .$$

Moreover, the TLS residual satisfies

$$(13) \quad \|(A, b) - (\tilde{A}, \tilde{b})\|_F^2 = -\lambda_I .$$

*Proof.* Define  $\tilde{r} \equiv b - \tilde{b} = b - \tilde{A}x$ . We characterize the solution to (7) by setting the partial derivatives of the Lagrangean  $\tilde{\mathcal{L}}$  (8) to zero. Differentiation with respect to the entries in  $\tilde{A}$  yields

$$\tilde{A} - A - \tilde{r}x^T = 0 ,$$

and differentiation with respect to the entries in  $x$  yields

$$(14) \quad -\tilde{A}^T \tilde{r} + \mu L^T L x = 0$$

or

$$(\tilde{A}^T \tilde{A} + \mu L^T L) x = \tilde{A}^T b .$$

Since  $A = \tilde{A} - \tilde{r}x^T$  and  $\tilde{A}^T \tilde{r} = \mu L^T L x$ , we see that

$$\begin{aligned} A^T A &= (\tilde{A}^T - x\tilde{r}^T)(\tilde{A} - \tilde{r}x^T) \\ &= \tilde{A}^T \tilde{A} - \mu x x^T L^T L + x\tilde{r}^T \tilde{r}x^T - \mu L^T L x x^T \end{aligned}$$

and

$$\tilde{A}^T b = A^T b + x\tilde{r}^T b .$$

Gathering terms, we arrive at (9), with

$$\begin{aligned} \lambda_I &= \mu \delta^2 - \|\tilde{r}\|_2^2 \|x\|_2^2 - \tilde{r}^T b , \\ \lambda_L &= \mu (1 + \|x\|_2^2) . \end{aligned}$$

To simplify the expression for  $\lambda_I$ , we first rewrite  $\tilde{r}$  as

$$\tilde{r} = b - \tilde{A}x = b - (A + \tilde{r}x^T)x = b - Ax - \tilde{r}x^T x$$

from which we obtain the relation

$$(15) \quad \tilde{r} (1 + \|x\|_2^2) = b - Ax$$

and therefore  $(1 + \|x\|_2^2)\|\tilde{r}\|_2^2 = (1 + \|x\|_2^2)^{-1}\|b - Ax\|_2^2$ . In (14), multiplication from the left by  $x^T$  leads to

$$(16) \quad \mu = \frac{x^T \tilde{A}^T \tilde{r}}{x^T L^T L x} = \frac{b^T \tilde{r} - \|\tilde{r}\|_2^2}{\delta^2} .$$

Inserting (15) and (16) into the above expression for  $\lambda_I$ , we obtain (10). Equation (12) is proved by multiplying  $\lambda_L$  by  $\delta^2$  and inserting (15) and (16). Finally, to prove the expression for the TLS residual, we use the relation

$$(A, b) - (\tilde{A}, \tilde{b}) = (A - \tilde{A}, b - \tilde{b}) = (-\tilde{r} x^T, \tilde{r}) = -\tilde{r} \begin{pmatrix} x \\ -1 \end{pmatrix}^T.$$

Taking the Frobenius norm and using (15) we obtain (13).  $\square$

Below, we discuss the implications of this theorem for both the standard-form case  $L = I_n$  (the identity matrix) and the general-form case  $L \neq I_n$ , both of which are important in applications.

**2.1. The Standard-Form Case.** In the standard-form case, Eq. (9) simplifies to

$$(17) \quad (A^T A + \lambda_{IL} I_n) x = A^T b$$

with  $\lambda_{IL} = \lambda_I + \lambda_L$ . The standard-form R-TLS solution  $\bar{x}_\delta$  to (7) has the same form as the standard-form Tikhonov solution  $x_\delta$  to (5). The two solutions have the following close relationship.

**THEOREM 2.2.** *Let  $L = I$  and let  $\bar{\sigma}_{n+1}$  denote the smallest singular value of  $(A, b)$ . Then for a given value of  $\delta$ , the solutions  $\bar{x}_\delta$  and  $x_\delta$  are related as follows:*

$\delta$	solutions	$\lambda_{IL}$
$\delta < \ x_{LS}\ _2$	$\bar{x}_\delta = x_\delta$	$\lambda_{IL} > 0$
$\delta = \ x_{LS}\ _2$	$\bar{x}_\delta = x_\delta = x_{LS}$	$\lambda_{IL} = 0$
$\ x_{LS}\ _2 < \delta < \ x_{TLS}\ _2$	$\bar{x}_\delta \neq x_\delta = x_{LS}$	$0 > \lambda_{IL} > -\bar{\sigma}_{n+1}^2$
$\delta \geq \ x_{TLS}\ _2$	$\bar{x}_\delta = x_{TLS}, x_\delta = x_{LS}$	$\lambda_{IL} = -\bar{\sigma}_{n+1}^2$

*Proof.* We need only determine the sign of  $\lambda_{IL}$  as a function of  $\delta$ , and for this we use the fact that  $\lambda_{IL}$  and its corresponding  $x$ -value are a solution to the Lagrangean formulation (6) of the Tikhonov regularization problem (5). The Lagrange multiplier  $\lambda$  for this problem is positive if  $\delta < \|x_{LS}\|_2$ , since the minimum value of the least squares residual can be decreased by relaxing the constraint, and is zero for  $\delta = \|x_{LS}\|_2$ . The multiplier is negative for  $\delta > \|x_{LS}\|_2$ , meaning that we can decrease the residual norm by allowing  $\|x\|_2$  to be less than  $\delta$ . If  $\delta = \|x_{TLS}\|_2$  then  $\lambda = -\bar{\sigma}_{n+1}^2$ , where  $\bar{\sigma}_{n+1}$  is the smallest singular value of  $(A, b)$  (see [16, Theorem 2.7]), and therefore we have  $-\bar{\sigma}_{n+1}^2 < \lambda < 0$  in the third case and  $\lambda = -\bar{\sigma}_{n+1}^2$  in the fourth.  $\square$

We conclude that as long as  $\delta \leq \|x_{LS}\|_2$ , which is normally the case in regularization problems since  $\|x_{LS}\|_2$  is very large, then R-TLS produces solutions that are identical to the Tikhonov solutions. In other words, replacing the least squares residual with the TLS residual in the Tikhonov formulation has no effect when  $L = I_n$  and  $\delta \leq \|x_{LS}\|_2$ .

We remark that since  $\|x_{TLS}\|_2 \geq \|x_{LS}\|_2$  (see [16, Corollary 6.2]) there is usually a nontrivial set of “large” values of  $\delta$  for which the multiplier  $\lambda_{IL}$  is negative. The corresponding R-TLS solutions  $\bar{x}_\delta$  are distinctly different from the Tikhonov solutions, and can be expected to be even more dominated by errors than the least squares solution  $x_{LS}$ .

We illustrate Theorem 2.2 with an example: discretization of a Fredholm integral equation with the second derivative operator as kernel. The implementation is `deriv2` from [9], the size of the matrix  $A$  is  $64 \times 32$ , and both  $A$  and  $b$  are perturbed by Gaussian noise with zero mean and standard deviation  $10^{-5}$ . We have  $\bar{\sigma}_{n+1}^2 \approx 2.38 \cdot$




FIG. 1. Regularized solutions to (3) for various values of the regularization parameter  $\lambda$ ; see Table 1 for details.

TABLE 1  
The solutions shown in Fig. 1.

solution	$\lambda$	solution norm	symbol
$x_\lambda$	$10^{-5}$	0.289	solid line
$x_{\text{LS}}$	0	0.385	dashed line
$x_\lambda$	$-\frac{1}{2}\bar{\sigma}_{n+1}^2$	0.408	crosses
$x_{\text{TLS}}$	$-\bar{\sigma}_{n+1}^2$	0.443	dotted line
$x^{\text{exact}}$	-	0.288	circles

$10^{-9}$ . Figure 1 shows the solutions listed in Table 1. We see that both  $x_{\text{LS}}$  and  $x_{\text{TLS}}$  have large oscillations (the oscillations of  $x_{\text{TLS}}$  being larger than those of  $x_{\text{LS}}$ ), while the regularized LS solution  $x_\lambda$  corresponding to  $\lambda = 10^{-5}$  is much smoother. The regularized TLS solution, corresponding to  $\lambda = -\frac{1}{2}\bar{\sigma}_{n+1}^2 = -1.19 \cdot 10^{-9}$ , has large oscillations whose amplitude is between that of  $x_{\text{TLS}}$  and  $x_{\text{LS}}$ .

We conclude that if we use Tikhonov regularization with  $L = I$  in order to reduce the norm of the solution below  $\|x_{\text{LS}}\|_2$ , then the regularized TLS formulation produces the same solution as the regularized LS formulation. This is in contrast to the truncating approach, since we have shown in [4] that truncated TLS can be superior to truncated LS.

**2.2. The General-Form Case.** In many applications, it is necessary to choose a matrix  $L$  different from the identity matrix; these issues are discussed, e.g., in [10, §4.3]. In this case, the R-TLS solution  $\bar{x}_\delta$  is different from the Tikhonov solution whenever the residual  $b - Ax$  is different from zero, since both  $\lambda_I$  and  $\lambda_L$  are nonzero.

From Theorem 2.2 we notice that  $\lambda_L$  is always positive when  $\delta < \|x_{\text{TLS}}\|_2$ , because the Lagrange parameter  $\mu$  is positive for these values of  $\lambda$ . On the other hand,  $\lambda_I$  is always negative, and thus adds some *de-regularization* to the solution. Statistical aspects of a negative regularization parameter in Tikhonov's method are discussed in [13].

For a given  $\delta$ , there are usually several pairs of parameters  $\lambda_I$  and  $\lambda_L$ , and thus several solutions  $x$ , that satisfy relations (9)–(11), but only one of these solves the optimization problem (7). According to (13), this is the solution that corresponds to the smallest value of  $|\lambda_I|$ . The following relations hold.

**THEOREM 2.3.** *For a given value of  $\delta$ , the solution  $\bar{x}_\delta$  is related to the solution to the total least squares problem as follows:*

$\delta$	solution	$\lambda_I$	$\lambda_L$
$\delta < \ L x_{\text{TLS}}\ _2$	$\bar{x}_\delta \neq x_{\text{TLS}}$	$\lambda_I < 0 \quad \partial\lambda_I/\partial\delta > 0$	$\lambda_L > 0$
$\delta \geq \ L x_{\text{TLS}}\ _2$	$\bar{x}_\delta = x_{\text{TLS}}$	$\lambda_I = -\bar{\sigma}_{n+1}^2$	$\lambda_L = 0$

*Proof.* For  $\delta < \|L x_{\text{TLS}}\|_2$ , the inequality constraint is active and the Lagrange multiplier  $\mu$  is positive, since relaxing the constraint decreases the residual. Thus  $\lambda_L$  is positive. The residual (13) is monotonically decreasing as  $\delta$  increases, since optimal solutions for smaller values of  $\delta$  are candidate solutions for larger  $\delta$ s, so  $\lambda_I$  is monotonically increasing.

For  $\delta = \|L x_{\text{TLS}}\|_2$ , the Lagrange multiplier is zero, and the solution becomes the unconstrained minimizer  $x_{\text{TLS}}$ . The value  $-\bar{\sigma}_{n+1}^2$  follows from [5, Theorem 4.1]. The constraint is never again active for larger  $\delta$ , so the solution remains unchanged.  $\square$

We note that if the matrix  $\lambda_I I_n + \lambda_L L^T L$  is positive definite, then the R-TLS solution corresponds to a Tikhonov solution for which the expression  $\lambda \|L x\|_2$  in (3) is replaced with the norm  $(\lambda_I \|x\|_2^2 + \lambda_L \|L x\|_2^2)^{1/2}$ . If  $\lambda_I I_n + \lambda_L L^T L$  is indefinite or negative definite then there is no equivalent interpretation.

**3. Computational Aspects.** To compute the R-TLS solutions for  $L \neq I_n$ , we have found it most convenient to avoid explicit use of  $\delta$ ; instead we use  $\lambda_L$  as the free parameter, fixing its value and then computing the value of  $\lambda_I$  that satisfies (10) and is smallest in absolute value. The corresponding value of  $\delta$  can then easily be computed from the relation (12).

We now discuss how to solve (9) efficiently for many values of  $\lambda_I$  and  $\lambda_L$ . First, we note that the equation is equivalent to the augmented system

$$(18) \quad \begin{pmatrix} I_m & 0 & A \\ 0 & I_p & \lambda_L^{1/2} L \\ A^T & \lambda_L^{1/2} L^T & -\lambda_I I_n \end{pmatrix} \begin{pmatrix} r \\ s \\ x \end{pmatrix} = \begin{pmatrix} b \\ 0 \\ 0 \end{pmatrix},$$

and our algorithm is based on this formulation.

We assume that the matrix  $L$  is a banded matrix, which is often the case when  $L$  represents a derivative operator. The key to efficiency is then to reduce  $A$  to  $n \times n$  bidiagonal form  $B$  by means of orthogonal transformations:  $H^T A K = B$ . The orthogonal right-transformations should also be applied to  $L$ , and simultaneously we should apply orthogonal transformations to  $L$  from the left in order to maintain its banded form. It is convenient to use sequences of Givens transformations to form  $J$ ,  $H$  and  $K$ , since this gives us the most freedom to retain the banded form of  $C = J^T L K$ .

Once  $B$  and  $C$  have been computed, we can recast the augmented system in (18) in the following form

$$(19) \quad \begin{pmatrix} I_n & 0 & B \\ 0 & I_p & \lambda_L^{1/2} C \\ B^T & \lambda_L^{1/2} C^T & -\lambda_I I_n \end{pmatrix} \begin{pmatrix} H^T r \\ J^T s \\ K^T x \end{pmatrix} = \begin{pmatrix} H^T b \\ 0 \\ 0 \end{pmatrix}.$$

Since  $\lambda_I$  changes more frequently than  $\lambda_L$  in our approach, we will now use Givens rotations to annihilate  $\lambda_L^{1/2} C$  using  $B$  by means of Elden's algorithm [1, Section 5.3.4], which can be represented as

$$\begin{pmatrix} B \\ \lambda_L^{1/2} C \end{pmatrix} = G \begin{pmatrix} \hat{B} \\ 0 \end{pmatrix} = \begin{pmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{pmatrix} \begin{pmatrix} \hat{B} \\ 0 \end{pmatrix}.$$






FIG. 2. Plots of the relative errors in the Tikhonov solutions (dashed lines) and R-TLS solutions (solid lines) versus  $\lambda_L$  for four values of the noise level.




FIG. 3. Plots of the "optimal" Tikhonov solutions (dashed lines) and R-TLS solutions (solid lines) for four values of the noise level. Also shown is the exact solution (dotted lines).

Figure 2 shows the relative errors  $\|x^* - x_\delta\|_2/\|x^*\|_2$  and  $\|x^* - \bar{x}_\delta\|_2/\|x^*\|_2$  in the Tikhonov and R-TLS solutions, respectively, for four values of the noise level:

$$\sigma = 0.03, 0.1, 0.3, 1.0.$$

We see that for small values of  $\sigma$  the two methods lead to almost the same minimum relative error, for almost the same value of  $\lambda_L$ . However, for larger value of  $\sigma$ , the minimum relative error for the R-TLS method is clearly smaller than that for Tikhonov's method, and it occurs for a larger value of  $\lambda_L$ . This shows the potential advantage of the R-TLS method, provided, of course, that a good estimate of the optimal regularization parameter can be found. This topic is outside the scope of the current paper.

In Fig. 3 we have plotted the "optimal" Tikhonov and R-TLS solutions, defined as the solutions that correspond to the minima of the curves in Fig. 2. In addition, we have plotted the exact solution  $x^*$ . Clearly, the addition of the term  $\lambda_I I_n$  in (9) introduces a non-constant component in the right part of the plot of the regularized solution, and it is precisely this component that improves the R-TLS error, compared to the Tikhonov error.

**Conclusion.** We have proved that the R-TLS solution to the regularized form (7) of the TLS problem is closely related to the Tikhonov solution to (3). In the standard form case,  $L = I_n$ , the two solutions are identical, showing that Tikhonov regularization in this case is also suited for problems with a perturbed coefficient matrix. For general problems with  $L \neq I_n$ , the R-TLS solution is different from the corresponding Tikhonov solution in that a deregularizing term  $\lambda_I I_n$  with a negative parameter  $\lambda_I$  is added to the formulation in (4).

#### REFERENCES

- [1] Å. BJÖRCK, *Numerical Methods for Least Squares Problems*, SIAM, Philadelphia, 1996.
- [2] H. W. ENGL, *Regularization methods for the stable solution of inverse problems*, *Surv. Math. Ind.* **3** (1993), 71–143.
- [3] H. W. ENGL, M. HANKE AND A. NEUBAUER, *Regularization of Inverse Problems*, Kluwer, Dordrecht, 1996.
- [4] R. D. FIERRO, G. H. GOLUB, P. C. HANSEN, D. P. O'LEARY, *Regularization by Truncated Total Least Squares*, Report UNIC-93-14, December 1993 (20 pages); submitted to *SIAM J. Sci. Comput.*
- [5] G. H. GOLUB AND C. F. VAN LOAN, *An analysis of the total least squares problem*, *SIAM J. Numer. Anal.* **17** (1980), 883–893.
- [6] G. H. GOLUB AND C. F. VAN LOAN, *Matrix Computations*, 3. Ed., Johns Hopkins University Press, Baltimore, 1996.
- [7] C. W. GROETSCH, *Inverse Problems in the Mathematical Sciences*, Vieweg, Wiesbaden, 1993.
- [8] M. HANKE AND P. C. HANSEN, *Regularization methods for large-scale problems*, *Surv. Math. Ind.* **3** (1993), 253–315.
- [9] P. C. HANSEN, *Regularization tools, a Matlab package for analysis of discrete regularization problems*, *Numerical Algorithms* **6** (1994), 1–35.
- [10] P. C. HANSEN, *Rank-Deficient and Discrete Ill-Posed Problems*, SIAM, Philadelphia, 1997.
- [11] P. C. HANSEN AND D. P. O'LEARY, *The use of the L-curve in the regularization of discrete ill-posed problems*, *SIAM J. Sci. Comput.*, **14** (1993), pp. 1487–1503.
- [12] P. C. HANSEN AND D. P. O'LEARY, *Regularization algorithms based on total least squares*; in S. Van Huffel (Ed.), *Recent Advances in Total Least Squares Techniques and Errors-in-Variables Modeling*, SIAM, Philadelphia, 1996; pp. 127–137.
- [13] T. A. HUA AND R. F. GUNST, *Generalized ridge regression: a note on negative ridge parameters*, *Commun. Statist. – Theor. Meth.*, **12** (1983), pp. 37–45.
- [14] G. W. STEWART, *On the invariance of perturbed null vectors under column scaling*, *Numer. Math.*, **44** (1984), pp. 61–65.

- [15] A. N. TIKHONOV AND V. Y. ARSENIN, *Solutions of Ill-Posed Problems*, Wiley, New York, 1977.
- [16] S. VAN HUFFEL AND J. VANDEWALLE, *The Total Least Squares Problem – Computational Aspects and Analysis*, SIAM, Philadelphia, 1991.
- [17] J. M. VARAH, *On the numerical solution of ill-conditioned linear systems with applications to ill-posed problems*, SIAM J. Numer. Anal., 10 (1973), pp. 257–267.