

## REFERENCES

- Aguayo, L. G. (1990). Ethanol potentiates the GABAA-activated Cl<sup>-</sup> current in mouse hippocampal and cortical neurons. *Eur J Pharmacol*, *187*(1), 127-130.
- Alex, K. D., Yavarian, G. J., McFarlane, H. G., Pluto, C. P., & Pehek, E. A. (2005). Modulation of dopamine release by striatal 5-HT<sub>2C</sub> receptors. *Synapse*, *55*(4), 242-251. doi: 10.1002/syn.20109
- Ballenger, J. C., Goodwin, F. K., Major, L. F., & Brown, G. L. (1979). Alcohol and central serotonin metabolism in man. *Arch Gen Psychiatry*, *36*(2), 224-227.
- Bardo, M. T., Donohew, R. L., & Harrington, N. G. (1996). Psychobiology of novelty seeking and drug seeking behavior. *Behav Brain Res*, *77*(1-2), 23-43.
- Begleiter, H., & Porjesz, B. (1988). Potential biological markers in individuals at high risk for developing alcoholism. *Alcohol Clin Exp Res*, *12*(4), 488-493.
- Berger, B., Gaspar, P., & Verney, C. (1991). Dopaminergic innervation of the cerebral cortex: unexpected differences between rodents and primates. *Trends Neurosci*, *14*(1), 21-27.
- Bhaskar, L. V., Thangaraj, K., Wasnik, S., Singh, L., & Raghavendra Rao, V. (2012). Dopamine transporter (DAT1) VNTR polymorphism and alcoholism in two culturally different populations of south India. *Am J Addict*, *21*(4), 343-347. doi: 10.1111/j.1521-0391.2012.00244.x
- Blanchard, B. A., Steindorf, S., Wang, S., & Glick, S. D. (1993). Sex differences in ethanol-induced dopamine release in nucleus accumbens and in ethanol consumption in rats. *Alcohol Clin Exp Res*, *17*(5), 968-973.
- Blednov, Y. A., & Harris, R. A. (2008). Metabotropic glutamate receptor 5 (mGluR5) regulation of ethanol sedation, dependence and consumption: relationship to acamprosate actions. *Int J Neuropsychopharmacol*, *11*(6), 775-793. doi: S1461145708008584 [pii] 10.1017/S1461145708008584
- Blomqvist, O., Gelernter, J., & Kranzler, H. R. (2000). Family-based study of DRD2 alleles in alcohol and drug dependence. *Am J Med Genet*, *96*(5), 659-664. doi: 10.1002/1096-8628(20001009)96:5<659::AID-AJMG12>3.0.CO;2-G [pii]
- Blum, K., Noble, E. P., Sheridan, P. J., Finley, O., Montgomery, A., Ritchie, T., . . . Sheffield, D. (1991). Association of the A1 allele of the D2 dopamine receptor gene with severe alcoholism. *Alcohol*, *8*(5), 409-416.

- Blum, K., Noble, E. P., Sheridan, P. J., Montgomery, A., Ritchie, T., Jagadeeswaran, P., . . . Cohn, J. B. (1990). Allelic association of human dopamine D2 receptor gene in alcoholism. *JAMA*, *263*(15), 2055-2060.
- Boyle, A. E., Smith, B. R., Spivak, K., & Amit, Z. (1994). Voluntary ethanol consumption in rats: the importance of the exposure paradigm in determining final intake outcome. *Behav Pharmacol*, *5*(4 And 5), 502-512.
- Brady, K. T., & Sonne, S. C. (1999). The role of stress in alcohol use, alcoholism treatment, and relapse. *Alcohol Res Health*, *23*(4), 263-271.
- Bronisch, T., & Wittchen, H.-U. (1992). Lifetime and 6-month prevalence of abuse and dependence of alcohol in the Munich follow-up study. *European Archives of Psychiatry and Clinical Neuroscience*, *241*(5), 273-282. doi: 10.1007/BF02195976
- Bullock, C. (1990). The biochemistry of alcohol metabolism — A brief review. *Biochemical Education*, *18*(2), 62-66. doi: 10.1016/0307-4412(90)90174-m
- Callebert, J., Esteve, J. M., Herve, P., Peoc'h, K., Tournois, C., Drouet, L., . . . Maroteaux, L. (2006). Evidence for a control of plasma serotonin levels by 5-hydroxytryptamine(2B) receptors in mice. *J Pharmacol Exp Ther*, *317*(2), 724-731. doi: 10.1124/jpet.105.098269
- Campbell, A. D., Kohl, R. R., & McBride, W. J. (1996). Serotonin-3 receptor and ethanol-stimulated somatodendritic dopamine release. *Alcohol*, *13*(6), 569-574. doi: S0741-8329(96)00069-9 [pii]
- Campbell, U. C., & Carroll, M. E. (2000). Acquisition of drug self-administration: environmental and pharmacological interventions. *Exp Clin Psychopharmacol*, *8*(3), 312-325.
- Carboni, E., & Vacca, C. (2003). Conditioned place preference. A simple method for investigating reinforcing properties in laboratory animals. *Methods Mol Med*, *79*, 481-498.
- Celentano, J. J., Gibbs, T. T., & Farb, D. H. (1988). Ethanol potentiates GABA- and glycine-induced chloride currents in chick spinal cord neurons. *Brain Res*, *455*(2), 377-380.
- Chaudhry, F. A., Edwards, R. H., & Fonnum, F. (2008). Vesicular neurotransmitter transporters as targets for endogenous and exogenous toxic substances. *Annu Rev Pharmacol Toxicol*, *48*, 277-301. doi: 10.1146/annurev.pharmtox.46.120604.141146
- Chen, N., Ferrer, J. V., Javitch, J. A., & Justice, J. B. (2000). Transport-dependent accessibility of a cytoplasmic loop cysteine in the human dopamine transporter. *J Biol Chem*, *275*(3),

1608-1614.

- Cloninger, C. R., Bohman, M., & Sigvardsson, S. (1981). Inheritance of alcohol abuse. Cross-fostering analysis of adopted men. *Arch Gen Psychiatry*, *38*(8), 861-868.
- Coull, J. T. (1998). Neural correlates of attention and arousal: insights from electrophysiology, functional neuroimaging and psychopharmacology. *Prog Neurobiol*, *55*(4), 343-361. doi: S0301-0082(98)00011-2 [pii]
- Cowen, M. S., & Lawrence, A. J. (1999). The role of opioid-dopamine interactions in the induction and maintenance of ethanol consumption. *Prog Neuropsychopharmacol Biol Psychiatry*, *23*(7), 1171-1212. doi: S0278-5846(99)00060-3 [pii]
- Crabbe, J. C., Spence, S. E., Brown, L. L., & Metten, P. (2011). Alcohol preference drinking in a mouse line selectively bred for high drinking in the dark. *Alcohol*, *45*(5), 427-440. doi: S0741-8329(10)00219-3 [pii]
- 10.1016/j.alcohol.2010.12.001
- Dahlström, A., & Fuxe, K. (1964). Localization of monoamines in the lower brain stem. *Experientia*, *20*(7), 398-399.
- Daoust, M., Chretien, P., Moore, N., Saligaut, C., Lhuintre, J. P., & Boismare, F. (1985). Isolation and striatal (3H) serotonin uptake: role in the voluntary intake of ethanol by rats. *Pharmacol Biochem Behav*, *22*(2), 205-208. doi: 0091-3057(85)90378-8 [pii]
- DHHS. (2000). *Tenth special report to congress on alcohol and health from the secretary of human services (DHHS Publication No. 00-1583)*. Washington, DC: U.S. Government Printing Office.
- Di Chiara, G. (1995). The role of dopamine in drug abuse viewed from the perspective of its role in motivation. *Drug Alcohol Depend*, *38*(2), 95-137. doi: 037687169501118I [pii]
- Diamond, I., & Gordon, A. S. (1997). Cellular and molecular neuroscience of alcoholism. *Physiol Rev*, *77*(1), 1-20.
- Diana, M., Pistis, M., Carboni, S., Gessa, G. L., & Rossetti, Z. L. (1993). Profound decrement of mesolimbic dopaminergic neuronal activity during ethanol withdrawal syndrome in rats: electrophysiological and biochemical evidence. *Proc Natl Acad Sci U S A*, *90*(17), 7966-7969.
- Dobashi, I., Inada, T., & Hadano, K. (1997). Alcoholism and gene polymorphisms related to central dopaminergic transmission in the Japanese population. *Psychiatr Genet*, *7*(2), 87-

91.

- Du, Y., Nie, Y., Li, Y., & Wan, Y. J. (2011). The association between the SLC6A3 VNTR 9-repeat allele and alcoholism-a meta-analysis. *Alcohol Clin Exp Res*, *35*(9), 1625-1634. doi: 10.1111/j.1530-0277.2011.01509.x
- Ducci, F., & Goldman, D. (2008). Genetic approaches to addiction: genes and alcohol. *Addiction*, *103*(9), 1414-1428. doi: 10.1111/j.1360-0443.2008.02203.x
- Edenberg, H. J., Koller, D. L., Xuei, X., Wetherill, L., McClintick, J. N., Almasy, L., . . . Foroud, T. (2010). Genome-wide association study of alcohol dependence implicates a region on chromosome 11. *Alcohol Clin Exp Res*, *34*(5), 840-852. doi: 10.1111/j.1530-0277.2010.01156.x
- Ehlers, C. L., & Schuckit, M. A. (1990). EEG fast frequency activity in the sons of alcoholics. *Biol Psychiatry*, *27*(6), 631-641.
- Eiden, L. E., Schäfer, M. K., Weihe, E., & Schütz, B. (2004). The vesicular amine transporter family (SLC18): amine/proton antiporters required for vesicular accumulation and regulated exocytotic secretion of monoamines and acetylcholine. *Pflugers Arch*, *447*(5), 636-640. doi: 10.1007/s00424-003-1100-5
- Eiden, L. E., & Weihe, E. (2011). VMAT2: a dynamic regulator of brain monoaminergic neuronal function interacting with drugs of abuse. *Ann N Y Acad Sci*, *1216*, 86-98. doi: 10.1111/j.1749-6632.2010.05906.x
- Elsworth, J. D., & Roth, R. H. (1997). Dopamine synthesis, uptake, metabolism, and receptors: relevance to gene therapy of Parkinson's disease. *Exp Neurol*, *144*(1), 4-9. doi: S0014-4886(96)96379-7 [pii] 10.1006/exnr.1996.6379
- Engleman, E. A., Keen, E. J., Tilford, S. S., Thielen, R. J., & Morzorati, S. L. (2011). Ethanol drinking reduces extracellular dopamine levels in the posterior ventral tegmental area of nondependent alcohol-preferring rats. *Alcohol*, *45*(6), 549-557. doi: S0741-8329(11)00358-2 [pii] 10.1016/j.alcohol.2011.02.304
- Enoch, M. A. (2012). The influence of gene-environment interactions on the development of alcoholism and drug dependence. *Curr Psychiatry Rep*, *14*(2), 150-158. doi: 10.1007/s11920-011-0252-9

- Erwin, V. G., & Deitrich, R. A. (1996). Genetic selection and characterization of mouse lines for acute functional tolerance to ethanol. *J Pharmacol Exp Ther*, *279*(3), 1310-1317.
- Fadda, F., Argiolas, A., Melis, M. R., Serra, G., & Gessa, G. L. (1980). Differential effect of acute and chronic ethanol on dopamine metabolism in frontal cortex, caudate nucleus and substantia nigra. *Life Sci*, *27*(11), 979-986.
- Fadda, F., & Rossetti, Z. L. (1998). Chronic ethanol consumption: from neuroadaptation to neurodegeneration. *Prog Neurobiol*, *56*(4), 385-431. doi: S0301-0082(98)00032-X [pii]
- Files, F. J., Lewis, R. S., & Samson, H. H. (1994). Effects of continuous versus limited access to ethanol on ethanol self-administration. *Alcohol*, *11*(6), 523-531. doi: 0741-8329(94)90079-5 [pii]
- Finckh, U., Rommelspacher, H., Kuhn, S., Dufeu, P., Otto, G., Heinz, A., . . . Rolfs, A. (1997). Influence of the dopamine D2 receptor (DRD2) genotype on neuroadaptive effects of alcohol and the clinical outcome of alcoholism. *Pharmacogenetics*, *7*(4), 271-281.
- Finckh, U., von Widdern, O., Giraldo-Velasquez, M., Podschus, J., Dufeu, P., Sander, T., . . . Rolfs, A. (1996). No association of the structural dopamine D2 receptor (DRD2) variant 311Cys with alcoholism. *Alcohol Clin Exp Res*, *20*(3), 528-532.
- Finn, P. R., & Pihl, R. O. (1988). Risk for alcoholism: a comparison between two different groups of sons of alcoholics on cardiovascular reactivity and sensitivity to alcohol. *Alcohol Clin Exp Res*, *12*(6), 742-747.
- Fitzpatrick, P. F. (1999). Tetrahydropterin-dependent amino acid hydroxylases. *Annu Rev Biochem*, *68*, 355-381. doi: 10.1146/annurev.biochem.68.1.355
- Fon, E. A., Pothos, E. N., Sun, B. C., Killeen, N., Sulzer, D., & Edwards, R. H. (1997). Vesicular transport regulates monoamine storage and release but is not essential for amphetamine action. *Neuron*, *19*(6), 1271-1283. doi: S0896-6273(00)80418-3 [pii]
- Froehlich, J. C. (1997). Opioid peptides. *Alcohol Health Res World*, *21*(2), 132-136.
- Gabrielli, W. F., Jr., Mednick, S. A., Volavka, J., Pollock, V. E., Schulsinger, F., & Itil, T. M. (1982). Electroencephalograms in children of alcoholic fathers. *Psychophysiology*, *19*(4), 404-407.
- Gainetdinov, R. R., Jones, S. R., Fumagalli, F., Wightman, R. M., & Caron, M. G. (1998). Re-evaluation of the role of the dopamine transporter in dopamine system homeostasis. *Brain Res Brain Res Rev*, *26*(2-3), 148-153.

- Gelernter, J., & Kranzler, H. (1999). D2 dopamine receptor gene (DRD2) allele and haplotype frequencies in alcohol dependent and control subjects: no association with phenotype or severity of phenotype. *Neuropsychopharmacology*, *20*(6), 640-649. doi: 10.1016/S0893-133X(98)00110-9
- Gelernter, J., O'Malley, S., Risch, N., Kranzler, H. R., Krystal, J., Merikangas, K., . . . Kidd, K. K. (1991). No association between an allele at the D2 dopamine receptor gene (DRD2) and alcoholism. *JAMA*, *266*(13), 1801-1807.
- George, S. R., Fan, T., Ng, G. Y., Jung, S. Y., O'Dowd, B. F., & Naranjo, C. A. (1995). Low endogenous dopamine function in brain predisposes to high alcohol preference and consumption: reversal by increasing synaptic dopamine. *J Pharmacol Exp Ther*, *273*(1), 373-379.
- Gianoulakis, C. (1989). The effect of ethanol on the biosynthesis and regulation of opioid peptides. *Experientia*, *45*(5), 428-435.
- Girault J, G. P. (2004). THE neurobiology of dopamine signaling. *Archives of Neurology*, *61*(5), 641-644. doi: 10.1001/archneur.61.5.641
- Giros, B., Jaber, M., Jones, S. R., Wightman, R. M., & Caron, M. G. (1996). Hyperlocomotion and indifference to cocaine and amphetamine in mice lacking the dopamine transporter. *Nature*, *379*(6566), 606-612. doi: 10.1038/379606a0
- Goldman, D., Oroszi, G., & Ducci, F. (2005). The genetics of addictions: uncovering the genes. *Nat Rev Genet*, *6*(7), 521-532. doi: 10.1038/nrg1635
- Goldman, D., Urbanek, M., Guenther, D., Robin, R., & Long, J. C. (1997). Linkage and association of a functional DRD2 variant [Ser311Cys] and DRD2 markers to alcoholism, substance abuse and schizophrenia in Southwestern American Indians. *Am J Med Genet*, *74*(4), 386-394.
- Goldman, D., Urbanek, M., Guenther, D., Robin, R., & Long, J. C. (1998). A functionally deficient DRD2 variant [Ser311Cys] is not linked to alcoholism and substance abuse. *Alcohol*, *16*(1), 47-52.
- Gonzales, R. A., & Weiss, F. (1998). Suppression of ethanol-reinforced behavior by naltrexone is associated with attenuation of the ethanol-induced increase in dialysate dopamine levels in the nucleus accumbens. *J Neurosci*, *18*(24), 10663-10671.
- Grant, K. A. (1994). Emerging neurochemical concepts in the actions of ethanol at ligand-gated

- ion channels. *Behav Pharmacol*, 5(4 And 5), 383-404.
- Greengard, P. (2001). The neurobiology of slow synaptic transmission. *Science*, 294(5544), 1024-1030. doi: 10.1126/science.294.5544.1024
- Grobin, A. C., Matthews, D. B., Devaud, L. L., & Morrow, A. L. (1998). The role of GABA[sub A] receptors in the acute and chronic effects of ethanol. *Psychopharmacology*, 139(1/2), 2.
- Guillot, T. S., & Miller, G. W. (2009). Protective actions of the vesicular monoamine transporter 2 (VMAT2) in monoaminergic neurons. *Mol Neurobiol*, 39(2), 149-170. doi: 10.1007/s12035-009-8059-y
- Hall, F. S., Huang, S., Fong, G. W., Pert, A., & Linnoila, M. (1998). Effects of isolation-rearing on voluntary consumption of ethanol, sucrose and saccharin solutions in Fawn Hooded and Wistar rats. *Psychopharmacology (Berl)*, 139(3), 210-216.
- Hall, F. S., Sora, I., & Uhl, G. R. (2001). Ethanol consumption and reward are decreased in mu-opiate receptor knockout mice. *Psychopharmacology (Berl)*, 154(1), 43-49.
- Hall, F. S., Sora, I., & Uhl, G. R. (2003). Sex-dependent modulation of ethanol consumption in vesicular monoamine transporter 2 (VMAT2) and dopamine transporter (DAT) knockout mice. *Neuropsychopharmacology*, 28(4), 620-628. doi: 1300070 [pii] 10.1038/sj.npp.1300070
- Hall, F. S., Wilkinson, L. S., Humby, T., Inglis, W., Kendall, D. A., Marsden, C. A., & Robbins, T. W. (1998). Isolation Rearing in Rats: Pre- and Postsynaptic Changes in Striatal Dopaminergic Systems. *Pharmacology Biochemistry and Behavior*, 59(4), 859-872. doi: 10.1016/S0091-3057(97)00510-8
- Hannon, J., & Hoyer, D. (2008). Molecular biology of 5-HT receptors. *Behav Brain Res*, 195(1), 198-213. doi: S0166-4328(08)00152-6 [pii] 10.1016/j.bbr.2008.03.020
- Harada, S., Agarwal, D. P., Goedde, H. W., & Ishikawa, B. (1983). Aldehyde dehydrogenase isozyme variation and alcoholism in Japan. *Pharmacol Biochem Behav*, 18 Suppl 1, 151-153.
- Hasin, D. S., Stinson, F. S., Ogburn, E., & Grant, B. F. (2007). Prevalence, correlates, disability, and comorbidity of DSM-IV alcohol abuse and dependence in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen*

- Psychiatry*, 64(7), 830-842. doi: 10.1001/archpsyc.64.7.830
- Hayashi, T. (1952). A physiological study of epileptic seizures following cortical stimulation in animals and its application to human clinics. *Jpn J Physiol*, 3(1), 46-64.
- Hayes, D. J., & Greenshaw, A. J. (2011). 5-HT receptors and reward-related behaviour: a review. *Neurosci Biobehav Rev*, 35(6), 1419-1449. doi: S0149-7634(11)00048-0 [pii] 10.1016/j.neubiorev.2011.03.005
- Heidbreder, C., & De Witte, P. (1993). Ethanol differentially affects extracellular monoamines and GABA in the nucleus accumbens. *Pharmacol Biochem Behav*, 46(2), 477-481.
- Heinz, A., Ragan, P., Jones, D. W., Hommer, D., Williams, W., Knable, M. B., . . . Linnoila, M. (1998). Reduced central serotonin transporters in alcoholism. *Am J Psychiatry*, 155(11), 1544-1549.
- Herz, A. (1997). Endogenous opioid systems and alcohol addiction. *Psychopharmacology (Berl)*, 129(2), 99-111.
- Higley, J. D., Suomi, S. J., & Linnoila, M. (1996). A nonhuman primate model of type II excessive alcohol consumption? Part 1. Low cerebrospinal fluid 5-hydroxyindoleacetic acid concentrations and diminished social competence correlate with excessive alcohol consumption. *Alcohol Clin Exp Res*, 20(4), 629-642.
- Higuchi, S., Muramatsu, T., Murayama, M., & Hayashida, M. (1994). Association of structural polymorphism of the dopamine D2 receptor gene and alcoholism. *Biochem Biophys Res Commun*, 204(3), 1199-1205. doi: 10.1006/bbrc.1994.2590
- Ho, P. S., Shih, M. C., Ma, K. H., Huang, W. S., Ho, K. K., Yen, C. H., . . . Huang, S. Y. (2011). Availability of the serotonin transporter in patients with alcohol dependence. *World J Biol Psychiatry*, 12(2), 134-142. doi: 10.3109/15622975.2010.503813
- Hodge, C. W., Samson, H. H., & Chappelle, A. M. (1997). Alcohol self-administration: further examination of the role of dopamine receptors in the nucleus accumbens. *Alcohol Clin Exp Res*, 21(6), 1083-1091.
- Hoyer, D., Hannon, J. P., & Martin, G. R. (2002). Molecular, pharmacological and functional diversity of 5-HT receptors. *Pharmacol Biochem Behav*, 71(4), 533-554.
- Johnson, B. (2004). The biologic basis of alcohol dependence. *Advanced Studies in Nursing*, 2(2), 48-53.
- Johnson, C., Drgon, T., Walther, D., & Uhl, G. R. (2011). Genomic regions identified by



- overlapping clusters of nominally-positive SNPs from genome-wide studies of alcohol and illegal substance dependence. *PLoS One*, *6*(7), e19210. doi: 10.1371/journal.pone.0019210
- Jones, S. R., Gainetdinov, R. R., Hu, X. T., Cooper, D. C., Wightman, R. M., White, F. J., & Caron, M. G. (1999). Loss of autoreceptor functions in mice lacking the dopamine transporter. *Nat Neurosci*, *2*(7), 649-655. doi: 10.1038/10204
- Jones, S. R., Gainetdinov, R. R., Jaber, M., Giros, B., Wightman, R. M., & Caron, M. G. (1998). Profound neuronal plasticity in response to inactivation of the dopamine transporter. *Proc Natl Acad Sci U S A*, *95*(7), 4029-4034.
- Jones, S. R., Gainetdinov, R. R., Wightman, R. M., & Caron, M. G. (1998). Mechanisms of amphetamine action revealed in mice lacking the dopamine transporter. *J Neurosci*, *18*(6), 1979-1986.
- Kebabian, J. W., & Calne, D. B. (1979). Multiple receptors for dopamine. *Nature*, *277*(5692), 93-96.
- Kendler, K. S., Gardner, C., & Dick, D. M. (2011). Predicting alcohol consumption in adolescence from alcohol-specific and general externalizing genetic risk factors, key environmental exposures and their interaction. *Psychol Med*, *41*(7), 1507-1516. doi: S003329171000190X [pii] 10.1017/S003329171000190X
- Kendler, K. S., Schmitt, E., Aggen, S. H., & Prescott, C. A. (2008). Genetic and environmental influences on alcohol, caffeine, cannabis, and nicotine use from early adolescence to middle adulthood. *Arch Gen Psychiatry*, *65*(6), 674-682. doi: 65/6/674 [pii] 10.1001/archpsyc.65.6.674
- Kopin, I. J. (1985). Catecholamine metabolism: basic aspects and clinical significance. *Pharmacol Rev*, *37*(4), 333-364.
- Lachowicz, J. E., & Sibley, D. R. (1997). Molecular characteristics of mammalian dopamine receptors. *Pharmacol Toxicol*, *81*(3), 105-113.
- Latvala, A., Tuulio-Henriksson, A., Perala, J., Saarni, S. I., Aalto-Setälä, T., Aro, H., . . . Suvisaari, J. (2009). Prevalence and correlates of alcohol and other substance use disorders in young adulthood: A population-based study. *BMC Psychiatry*, *9*, 73. doi: 10.1186/1471-244x-9-73

- Le, A. D., Ko, J., Chow, S., & Quan, B. (1994). Alcohol consumption by C57BL/6, BALB/c, and DBA/2 mice in a limited access paradigm. *Pharmacol Biochem Behav*, *47*(2), 375-378.
- Le Strat, Y., Ramoz, N., Pickering, P., Burger, V., Boni, C., Aubin, H. J., . . . Gorwood, P. (2008). The 3' part of the dopamine transporter gene DAT1/SLC6A3 is associated with withdrawal seizures in patients with alcohol dependence. *Alcohol Clin Exp Res*, *32*(1), 27-35. doi: 10.1111/j.1530-0277.2007.00552.x
- LeMarquand, D., Pihl, R. O., & Benkelfat, C. (1994). Serotonin and alcohol intake, abuse, and dependence: clinical evidence. *Biol Psychiatry*, *36*(5), 326-337. doi: 0006-3223(94)90630-0 [pii]
- Lim, J. P., Zou, M. E., Janak, P. H., & Messing, R. O. (2012). Responses to ethanol in C57BL/6 versus C57BL/6 x 129 hybrid mice. *Brain Behav*, *2*(1), 22-31. doi: 10.1002/brb3.29
- Lin, Z., & Uhl, G. R. (2003). Human dopamine transporter gene variation: effects of protein coding variants V55A and V382A on expression and uptake activities. *Pharmacogenomics J*, *3*(3), 159-168. doi: 10.1038/sj.tpj.6500169
- Lin, Z., Walther, D., Yu, X. Y., Li, S., Drgon, T., & Uhl, G. R. (2005). SLC18A2 promoter haplotypes and identification of a novel protective factor against alcoholism. *Hum Mol Genet*, *14*(10), 1393-1404. doi: 10.1093/hmg/ddi148
- Linnoila, M., Virkkunen, M., Scheinin, M., Nuutila, A., Rimon, R., & Goodwin, F. K. (1983). Low cerebrospinal fluid 5-hydroxyindoleacetic acid concentration differentiates impulsive from nonimpulsive violent behavior. *Life Sci*, *33*(26), 2609-2614.
- Liu, J., Zhou, Z., Hodgkinson, C. A., Yuan, Q., Shen, P. H., Mulligan, C. J., . . . Enoch, M. A. (2011). Haplotype-based study of the association of alcohol-metabolizing genes with alcohol dependence in four independent populations. *Alcohol Clin Exp Res*, *35*(2), 304-316. doi: 10.1111/j.1530-0277.2010.01346.x
- Ljungberg, T., Apicella, P., & Schultz, W. (1992). Responses of monkey dopamine neurons during learning of behavioral reactions. *J Neurophysiol*, *67*(1), 145-163.
- Lovinger, D. M., White, G., & Weight, F. F. (1989). Ethanol inhibits NMDA-activated ion current in hippocampal neurons. *Science*, *243*(4899), 1721-1724.
- Lucki, I. (1998). The spectrum of behaviors influenced by serotonin. *Biol Psychiatry*, *44*(3), 151-162.

- Lyness, W. H., & Smith, F. L. (1992). Influence of dopaminergic and serotonergic neurons on intravenous ethanol self-administration in the rat. *Pharmacol Biochem Behav*, *42*(1), 187-192.
- Mateo, Y., Budygin, E. A., John, C. E., Banks, M. L., & Jones, S. R. (2004). Voltammetric assessment of dopamine clearance in the absence of the dopamine transporter: no contribution of other transporters in core or shell of nucleus accumbens. *J Neurosci Methods*, *140*(1-2), 183-187. doi: 10.1016/j.jneumeth.2004.05.018
- Mayfield, R. D., Maiya, R., Keller, D., & Zahniser, N. R. (2001). Ethanol potentiates the function of the human dopamine transporter expressed in *Xenopus* oocytes. *Journal of Neurochemistry*, *79*(5), 1070-1079.
- McCormick, D. A. (1989). GABA as an inhibitory neurotransmitter in human cerebral cortex. *J Neurophysiol*, *62*(5), 1018-1027.
- McGue, M., Pickens, R. W., & Svikis, D. S. (1992). Sex and age effects on the inheritance of alcohol problems: a twin study. *J Abnorm Psychol*, *101*(1), 3-17.
- Mereu, G., & Gessa, G. L. (1985). Low doses of ethanol inhibit the firing of neurons in the substantia nigra, pars reticulata: a GABAergic effect? *Brain Res*, *360*(1-2), 325-330. doi: 0006-8993(85)91249-1 [pii]
- Mihic, S. J., & Harris, R. A. (1997). GABA and the GABAA receptor. *Alcohol Health Res World*, *21*(2), 127-131.
- Molinoff, P. B., & Axelrod, J. (1971). Biochemistry of catecholamines. *Annu Rev Biochem*, *40*, 465-500. doi: 10.1146/annurev.bi.40.070171.002341
- Morice, E., Denis, C., Giros, B., & Nosten-Bertrand, M. (2010). Evidence of long-term expression of behavioral sensitization to both cocaine and ethanol in dopamine transporter knockout mice. *Psychopharmacology (Berl)*, *208*(1), 57-66. doi: 10.1007/s00213-009-1707-0
- Moron, J. A., Brockington, A., Wise, R. A., Rocha, B. A., & Hope, B. T. (2002). Dopamine uptake through the norepinephrine transporter in brain regions with low levels of the dopamine transporter: evidence from knock-out mouse lines. *J Neurosci*, *22*(2), 389-395.
- Muramatsu, T., & Higuchi, S. (1995). Dopamine transporter gene polymorphism and alcoholism. *Biochem Biophys Res Commun*, *211*(1), 28-32. doi: 10.1006/bbrc.1995.1773
- Narboux-Neme, N., Sagne, C., Doly, S., Diaz, S. L., Martin, C. B., Angenard, G., . . . Mongeau,

- R. (2011). Severe serotonin depletion after conditional deletion of the vesicular monoamine transporter 2 gene in serotonin neurons: neural and behavioral consequences. *Neuropsychopharmacology*, *36*(12), 2538-2550. doi: 10.1038/npp.2011.142
- Neve, K. A., Seamans, J. K., & Trantham-Davidson, H. (2004). Dopamine receptor signaling. *J Recept Signal Transduct Res*, *24*(3), 165-205.
- Newlin, D. B. (1985). Offspring of alcoholics have enhanced antagonistic placebo response. *J Stud Alcohol*, *46*(6), 490-494.
- Newlin, D. B., & Thomson, J. B. (1991). Chronic tolerance and sensitization to alcohol in sons of alcoholics. *Alcohol Clin Exp Res*, *15*(3), 399-405.
- Nicola, S. M., Surmeier, D. J., & Malenka, R. C. (2000). Dopaminergic Modulation of Neuronal Excitability in the Striatum and Nucleus Accumbens. *Annual Review of Neuroscience*, *23*(1), 185-215. doi: doi:10.1146/annurev.neuro.23.1.185
- Nielsen, D. A., Goldman, D., Virkkunen, M., Tokola, R., Rawlings, R., & Linnoila, M. (1994). Suicidality and 5-hydroxyindoleacetic acid concentration associated with a tryptophan hydroxylase polymorphism. *Arch Gen Psychiatry*, *51*(1), 34-38.
- Noble, E. P., Blum, K., Ritchie, T., Montgomery, A., & Sheridan, P. J. (1991). Allelic association of the D2 dopamine receptor gene with receptor-binding characteristics in alcoholism. *Arch Gen Psychiatry*, *48*(7), 648-654.
- Olbrich, H. M., Maes, H., Valerius, G., Langosch, J. M., Gann, H., & Feige, B. (2002). Assessing cerebral dysfunction with probe-evoked potentials in a CNV task -- a study in alcoholics. *Clin Neurophysiol*, *113*(6), 815-825.
- Ono, Y., Manki, H., Yoshimura, K., Muramatsu, T., Mizushima, H., Higuchi, S., . . . Asai, M. (1997). Association between dopamine D4 receptor (D4DR) exon III polymorphism and novelty seeking in Japanese subjects. *Am J Med Genet*, *74*(5), 501-503.
- Parsian, A., & Zhang, Z. H. (1997). Human dopamine transporter gene polymorphism (VNTR) and alcoholism. *Am J Med Genet*, *74*(5), 480-482.
- Parsons, S. M. (2000). Transport mechanisms in acetylcholine and monoamine storage. *FASEB J*, *14*(15), 2423-2434. doi: 10.1096/fj.00-0203rev
- Partilla, J. S., Dempsey, A. G., Nagpal, A. S., Blough, B. E., Baumann, M. H., & Rothman, R. B. (2006). Interaction of amphetamines and related compounds at the vesicular monoamine transporter. *J Pharmacol Exp Ther*, *319*(1), 237-246. doi: 10.1124/jpet.106.103622

- Peris, J., Coleman-Hardee, M., Burry, J., & Pecins-Thompson, M. (1992). Selective changes in GABAergic transmission in substantia nigra and superior colliculus caused by ethanol and ethanol withdrawal. *Alcohol Clin Exp Res*, *16*(2), 311-319.
- Pickens, R. W., Svikis, D. S., McGue, M., Lykken, D. T., Heston, L. L., & Clayton, P. J. (1991). Heterogeneity in the inheritance of alcoholism. A study of male and female twins. *Arch Gen Psychiatry*, *48*(1), 19-28.
- Pollock, V. E., Gabrielli, W. F., Mednick, S. A., & Goodwin, D. W. (1988). EEG identification of subgroups of men at risk for alcoholism? *Psychiatry Res*, *26*(1), 101-104.
- Porrino, L. J., & Goldman-Rakic, P. S. (1982). Brainstem innervation of prefrontal and anterior cingulate cortex in the rhesus monkey revealed by retrograde transport of HRP. *J Comp Neurol*, *205*(1), 63-76. doi: 10.1002/cne.902050107
- Prescott, C. A., Aggen, S. H., & Kendler, K. S. (1999). Sex differences in the sources of genetic liability to alcohol abuse and dependence in a population-based sample of U.S. twins. *Alcohol Clin Exp Res*, *23*(7), 1136-1144.
- Prescott, C. A., & Kendler, K. S. (1999). Genetic and environmental contributions to alcohol abuse and dependence in a population-based sample of male twins. *Am J Psychiatry*, *156*(1), 34-40.
- Qiang, M., Denny, A. D., & Ticku, M. K. (2007). Chronic intermittent ethanol treatment selectively alters N-methyl-D-aspartate receptor subunit surface expression in cultured cortical neurons. *Mol Pharmacol*, *72*(1), 95-102. doi: mol.106.033043 [pii] 10.1124/mol.106.033043
- Rausch, J. L., Monteiro, M. G., & Schuckit, M. A. (1991). Platelet serotonin uptake in men with family histories of alcoholism. *Neuropsychopharmacology*, *4*(2), 83-86.
- Renson, J., Goodwin, F., Weissbach, H., & Udenfriend, S. (1961). Conversion of tryptophan to 5-hydroxytryptophan by phenylalanine hydroxylase. *Biochem Biophys Res Commun*, *6*, 20-23.
- Rhodes, J. S., Best, K., Belknap, J. K., Finn, D. A., & Crabbe, J. C. (2005). Evaluation of a simple model of ethanol drinking to intoxication in C57BL/6J mice. *Physiol Behav*, *84*(1), 53-63. doi: 10.1016/j.physbeh.2004.10.007
- Roberts, A. J., McDonald, J. S., Heyser, C. J., Kieffer, B. L., Matthes, H. W., Koob, G. F., & Gold, L. H. (2000). mu-Opioid receptor knockout mice do not self-administer alcohol. *J*

- Pharmacol Exp Ther*, 293(3), 1002-1008.
- Samochowiec, J., Kucharska-Mazur, J., Grzywacz, A., Jablonski, M., Rommelspacher, H., Samochowiec, A., . . . Pelka-Wysiecka, J. (2006). Family-based and case-control study of DRD2, DAT, 5HTT, COMT genes polymorphisms in alcohol dependence. *Neurosci Lett*, 410(1), 1-5. doi: 10.1016/j.neulet.2006.05.005
- Sander, T., Harms, H., Dufeu, P., Kuhn, S., Rommelspacher, H., & Schmidt, L. G. (1997). Dopamine D4 receptor exon III alleles and variation of novelty seeking in alcoholics. *Am J Med Genet*, 74(5), 483-487.
- Sander, T., Harms, H., Podschus, J., Finckh, U., Nickel, B., Rolfs, A., . . . Schmidt, L. G. (1997). Allelic association of a dopamine transporter gene polymorphism in alcohol dependence with withdrawal seizures or delirium. *Biol Psychiatry*, 41(3), 299-304.
- Savelieva, K. V., Caudle, W. M., Findlay, G. S., Caron, M. G., & Miller, G. W. (2002). Decreased ethanol preference and consumption in dopamine transporter female knock-out mice. *Alcohol Clin Exp Res*, 26(6), 758-764.
- Savelieva, K. V., Caudle, W. M., & Miller, G. W. (2006). Altered ethanol-associated behaviors in vesicular monoamine transporter heterozygote knockout mice. *Alcohol*, 40(2), 87-94. doi: 10.1016/j.alcohol.2006.09.030
- Schellekens, A. F., Franke, B., Ellenbroek, B., Cools, A., de Jong, C. A., Buitelaar, J. K., & Verkes, R. J. (2012). Reduced dopamine receptor sensitivity as an intermediate phenotype in alcohol dependence and the role of the COMT Val158Met and DRD2 Taq1A genotypes. *Arch Gen Psychiatry*, 69(4), 339-348. doi: 10.1001/archgenpsychiatry.2011.1335
- Schuckit, M. A. (1994). A clinical model of genetic influences in alcohol dependence. *J Stud Alcohol*, 55(1), 5-17.
- Schultz, W. (1997). Dopamine neurons and their role in reward mechanisms. *Curr Opin Neurobiol*, 7(2), 191-197.
- Schultz, W. (1999). The Reward Signal of Midbrain Dopamine Neurons. *News Physiol Sci*, 14, 249-255.
- Shen, Y. C., Fan, J. H., Edenberg, H. J., Li, T. K., Cui, Y. H., Wang, Y. F., . . . Xia, G. Y. (1997). Polymorphism of ADH and ALDH genes among four ethnic groups in China and effects upon the risk for alcoholism. *Alcohol Clin Exp Res*, 21(7), 1272-1277.

- Simpson, E. M., Linder, C. C., Sargent, E. E., Davisson, M. T., Mobraaten, L. E., & Sharp, J. J. (1997). Genetic variation among 129 substrains and its importance for targeted mutagenesis in mice. *Nat Genet*, *16*(1), 19-27. doi: 10.1038/ng0597-19
- Smith, A. D., & Weiss, F. (1999). Ethanol exposure differentially alters central monoamine neurotransmission in alcohol-preferring versus -nonpreferring rats. *J Pharmacol Exp Ther*, *288*(3), 1223-1228.
- Sora, I., Elmer, G., Funada, M., Pieper, J., Li, X. F., Hall, F. S., & Uhl, G. R. (2001). Mu opiate receptor gene dose effects on different morphine actions: evidence for differential in vivo mu receptor reserve. *Neuropsychopharmacology*, *25*(1), 41-54. doi: 10.1016/s0893-133x(00)00252-9
- Sora, I., Hall, F. S., Andrews, A. M., Itokawa, M., Li, X. F., Wei, H. B., . . . Uhl, G. R. (2001). Molecular mechanisms of cocaine reward: combined dopamine and serotonin transporter knockouts eliminate cocaine place preference. *Proc Natl Acad Sci U S A*, *98*(9), 5300-5305. doi: 10.1073/pnas.091039298
- Stamatoyannopoulos, G., Chen, S. H., & Fukui, M. (1975). Liver alcohol dehydrogenase in Japanese: high population frequency of atypical form and its possible role in alcohol sensitivity. *Am J Hum Genet*, *27*(6), 789-796.
- Stamford, J. A., Kruk, Z. L., & Millar, J. (1990). Striatal dopamine terminals release serotonin after 5-HTP pretreatment: in vivo voltammetric data. *Brain Res*, *515*(1-2), 173-180. doi: 0006-8993(90)90593-Z [pii]
- Suzdak, P. D., Schwartz, R. D., Skolnick, P., & Paul, S. M. (1986). Ethanol stimulates gamma-aminobutyric acid receptor-mediated chloride transport in rat brain synaptoneuroosomes. *Proc Natl Acad Sci U S A*, *83*(11), 4071-4075.
- Tabakoff, B., & Hoffman, P. L. (2000). Animal models in alcohol research. *Alcohol Res Health*, *24*(2), 77-84.
- Tordoff, M. G., & Bachmanov, A. A. (2002). Influence of test duration on the sensitivity of the two-bottle choice test. *Chem Senses*, *27*(9), 759-768.
- Torres, G. E., Gainetdinov, R. R., & Caron, M. G. (2003). Plasma membrane monoamine transporters: structure, regulation and function. *Nat Rev Neurosci*, *4*(1), 13-25. doi: 10.1038/nrn1008
- Tossman, U., Jonsson, G., & Ungerstedt, U. (1986). Regional distribution and extracellular

- levels of amino acids in rat central nervous system. *Acta Physiologica Scandinavica*, *127*(4), 533-545. doi: 10.1111/j.1748-1716.1986.tb07938.x
- Treutlein, J., Cichon, S., Ridinger, M., Wodarz, N., Soyka, M., Zill, P., . . . Rietschel, M. (2009). Genome-wide association study of alcohol dependence. *Arch Gen Psychiatry*, *66*(7), 773-784. doi: 10.1001/archgenpsychiatry.2009.83
- Tsai, G. E., & Coyle, J. T. (1998). The role of glutamatergic neurotransmission in the pathophysiology of alcoholism. *Annu Rev Med*, *49*, 173-184. doi: 10.1146/annurev.med.49.1.173
- Tsai, G. E., Ragan, P., Chang, R., Chen, S., Linnoila, V. M., & Coyle, J. T. (1998). Increased glutamatergic neurotransmission and oxidative stress after alcohol withdrawal. *Am J Psychiatry*, *155*(6), 726-732.
- Ueno, S., Nakamura, M., Mikami, M., Kondoh, K., Ishiguro, H., Arinami, T., . . . Tanabe, H. (1999). Identification of a novel polymorphism of the human dopamine transporter (DAT1) gene and the significant association with alcoholism. *Mol Psychiatry*, *4*(6), 552-557.
- Uhl, G., Blum, K., Noble, E., & Smith, S. (1993). Substance abuse vulnerability and D2 receptor genes. *Trends Neurosci*, *16*(3), 83-88.
- Uhl, G. R. (1992). Neurotransmitter transporters (plus): a promising new gene family. *Trends Neurosci*, *15*(7), 265-268.
- Ulm, R. R., Volpicelli, J. R., & Volpicelli, L. A. (1995). Opiates and alcohol self-administration in animals. *J Clin Psychiatry*, *56 Suppl 7*, 5-14.
- Vandenbergh, D. J., Thompson, M. D., Cook, E. H., Bendahhou, E., Nguyen, T., Krasowski, M. D., . . . Uhl, G. R. (2000). Human dopamine transporter gene: coding region conservation among normal, Tourette's disorder, alcohol dependence and attention-deficit hyperactivity disorder populations. *Mol Psychiatry*, *5*(3), 283-292.
- Wang, Y. M., Gainetdinov, R. R., Fumagalli, F., Xu, F., Jones, S. R., Bock, C. B., . . . Caron, M. G. (1997). Knockout of the vesicular monoamine transporter 2 gene results in neonatal death and supersensitivity to cocaine and amphetamine. *Neuron*, *19*(6), 1285-1296.
- Wauquier, A. (1980). The pharmacology of catecholamine involvement in the neural mechanisms of reward. *Acta Neurobiol Exp (Wars)*, *40*(3), 665-686.
- Werkman, T. R., Glennon, J. C., Wadman, W. J., & McCreary, A. C. (2006). Dopamine receptor



- pharmacology: interactions with serotonin receptors and significance for the aetiology and treatment of schizophrenia. *CNS Neurol Disord Drug Targets*, *5(1)*, 3-23.
- Westlund, K. N., Denney, R. M., Kochersperger, L. M., Rose, R. M., & Abell, C. W. (1985). Distinct monoamine oxidase A and B populations in primate brain. *Science*, *230(4722)*, 181-183.
- Williams, S. M., & Goldman-Rakic, P. S. (1998). Widespread origin of the primate mesofrontal dopamine system. *Cereb Cortex*, *8(4)*, 321-345.
- Woodward, J. J. (1999). Ionotropic glutamate receptors as sites of action for ethanol in the brain. *Neurochem Int*, *35(2)*, 107-113. doi: S0197-0186(99)00052-2 [pii]
- Wozniak, K. M., Pert, A., & Linnoila, M. (1990). Antagonism of 5-HT<sub>3</sub> receptors attenuates the effects of ethanol on extracellular dopamine. *Eur J Pharmacol*, *187(2)*, 287-289. doi: 0014-2999(90)90015-X [pii]
- Xiong, N., Huang, J., Chen, C., Zhao, Y., Zhang, Z., Jia, M., . . . Wang, T. (2012). D1-3-n-butylphthalide, a natural antioxidant, protects dopamine neurons in rotenone models for Parkinson's disease. *Neurobiol Aging*, *33(8)*, 1777-1791. doi: 10.1016/j.neurobiolaging.2011.03.007
- Xiong, N., Xiong, J., Khare, G., Chen, C., Huang, J., Zhao, Y., . . . Wang, T. (2011). Edaravone guards dopamine neurons in a rotenone model for Parkinson's disease. *PLoS One*, *6(6)*, e20677. doi: 10.1371/journal.pone.0020677
- Yamamoto, H., Kamegaya, E., Hagino, Y., Imai, K., Fujikawa, A., Tamura, K., . . . Sora, I. (2007). Genetic deletion of vesicular monoamine transporter-2 (VMAT2) reduces dopamine transporter activity in mesencephalic neurons in primary culture. *Neurochem Int*, *51(2-4)*, 237-244. doi: 10.1016/j.neuint.2007.06.022
- Yan, Q. S., Reith, M. E., Jobe, P. C., & Dailey, J. W. (1996). Focal ethanol elevates extracellular dopamine and serotonin concentrations in the rat ventral tegmental area. *Eur J Pharmacol*, *301(1-3)*, 49-57.
- Yasumoto, S., Tamura, K., Karasawa, J., Hasegawa, R., Ikeda, K., Yamamoto, T., & Yamamoto, H. (2009). Inhibitory effect of selective serotonin reuptake inhibitors on the vesicular monoamine transporter 2. *Neurosci Lett*, *454(3)*, 229-232. doi: 10.1016/j.neulet.2009.03.049
- Yoneyama, N., Crabbe, J. C., Ford, M. M., Murillo, A., & Finn, D. A. (2008). Voluntary ethanol

consumption in 22 inbred mouse strains. *Alcohol*, 42(3), 149-160. doi: S0741-8329(08)00024-4 [pii]

10.1016/j.alcohol.2007.12.006

Yoshimoto, K., Yayama, K., Sorimachi, Y., Tani, J., Ogata, M., Nishimura, A., . . . Komura, S. (1996). Possibility of 5-HT<sub>3</sub> receptor involvement in alcohol dependence: a microdialysis study of nucleus accumbens dopamine and serotonin release in rats with chronic alcohol consumption. *Alcohol Clin Exp Res*, 20(9 Suppl), 311A-319A.

Zakhari, S. (2006). Overview: how is alcohol metabolized by the body? *Alcohol Res Health*, 29(4), 245-254.

Zhou, F. C., Lesch, K. P., & Murphy, D. L. (2002). Serotonin uptake into dopamine neurons via dopamine transporters: a compensatory alternative. *Brain Res*, 942(1-2), 109-119. doi: S0006899302027099 [pii]