Supplemental Table 1. Stroke Volume, Cardiac Output, and Total Peripheral Resistance Responses to the Functional Sympatholysis Protocol.

	Baseline A	**CPT**	**Baseline B**	**Exercise**	**Exercise + CPT**
SV, mL (N=13)	83.22 ± 21.91	79.08 ± 22.01 #	85.99 ±19.06	85.06 ± 22.10	79.82 ± 23.61
Females (N=7)	77.49 ± 14.11	73.61 ± 16.08	77.97 ± 12.58	78.98 ± 18.55	75.42 ± 19.91
Males (N=6)	89.90 ± 28.53	85.47 ± 27.59	95.34 ± 22.07	92.16 ± 25.43	84.96 ± 28.34
Q, L/min (N=13)	5.01 ± 1.13	5.59 ± 1.40 #	5.13 ± 0.82	6.24 ± 1.31 #	5.99 ± 1.46 #
Females (N=7)	5.23 ± 1.20	5.71 ± 1.28	5.25 ± 0.87	6.29 ± 1.54	6.01 ± 1.62
Males (N=6)	4.76 ± 1.08	5.44 ± 1.63	4.98 ± 0.80	6.18 ± 1.12	5.97 ± 1.40
TPR, mmHg*s/mL (N=13)	1.21 ± 0.39	1.38 ± 0.49	1.11 ± 0.24	1.12 ± 0.45	1.44 ± 0.64 # $
Females (N=7)	1.15 ± 0.42	1.28 ± 0.53	1.04 ± 0.22	1.07 ± 0.47	1.28 ± 0.61
Males (N=6)	1.28 ± 0.37	1.50 ± 0.45	1.19 ± 0.25	1.16 ± 0.47	1.62 ± 0.68

Supplemental Table 1 Key: CPT; cold-pressor task; SV, stroke volume; Q, cardiac output; TPR, total peripheral resistance. * indicates significantly different ($P<0.05$) from males during the same condition. # indicates significantly different ($P<0.05$) from the baseline condition of that trial. $ indicates significant difference ($P<0.05$) between the Ex and Ex+CPT conditions.