

On The Number Of Unlabeled Bipartite Graphs

Abdullah Atmaca and A. Yavuz Oruç

The
Institute for
Systems
Research



A. JAMES CLARK
SCHOOL OF ENGINEERING

ISR develops, applies and teaches advanced methodologies of design and analysis to solve complex, hierarchical, heterogeneous and dynamic problems of engineering technology and systems for industry and government.

ISR is a permanent institute of the University of Maryland, within the A. James Clark School of Engineering. It is a graduated National Science Foundation Engineering Research Center.

On The Number Of Unlabeled Bipartite Graphs

Abdullah Atmaca* and A. Yavuz Oruç†

Abstract

Let I and O denote two sets of vertices, where $I \cap O = \Phi$, $|I| = n$, $|O| = r$, and $B_u(n, r)$ denote the set of unlabeled graphs whose edges connect vertices in I and O . It is shown that the following two-sided equality holds.

$$\frac{\binom{r+2^n-1}{r}}{n!} \leq |B_u(n, r)| \leq 2 \frac{\binom{r+2^n-1}{r}}{n!} \quad (1)$$

where $n < r$.

1 Introduction

This work was motivated in part by a counting problem that arises in the representation of calls in interconnection networks [1]. It has also been investigated in connection with the enumeration of unlabeled bipartite graphs and binary matrices[2]. Let (I, O, E) denote a graph with two disjoint sets of vertices, I , called *left vertices* and a set of vertices, O , called *right vertices*, where each edge in E connects a left vertex with a right vertex. We let $n = |I|$, $r = |O|$, and refer to such a graph as an (n, r) -bipartite graph. Let $G_1 = (I, O, E_1)$ and $G_2 = (I, O, E_2)$ be two (n, r) -bipartite graphs, and $\alpha : I \rightarrow I$ and $\beta : O \rightarrow O$ be both bijections. The pair (α, β) is an isomorphism between G_1 and G_2 provided that $((\alpha(v_1), \beta(v_2)) \in E_2$ if and only if $(v_1, v_2) \in E_1$, $\forall v_1 \in I, \forall v_2 \in O$. It is easy to establish that this mapping induces an equivalence relation, and partitions the set of 2^{nr} (n, r) -bipartite graphs into equivalence classes. This equivalence relation captures the fact that the vertices in I and O are unlabeled, and so each class of (n, r) -bipartite graphs can be represented by any one of the graphs in that class without identifying the vertices in I and O . Let $B_u(n, r)$ denote any set of (n, r) -bipartite graphs that contains exactly one such graph from each of the equivalence classes of (n, r) -bipartite graphs induced by the isomorphism we defined. It is easy to see that determining $|B_u(n, r)|$ amounts to an enumeration of non-isomorphic (n, r) -bipartite graphs that will henceforth be referred to as unlabeled (n, r) -bipartite graphs.

In [2], Harrison used Pólya's counting theorem to obtain an expression to compute the number of non-equivalent $n \times r$ binary matrices. This expression contains a nested sum, in which one sum is carried over all partitions of n while the other is carried over all partitions of r , where the argument of the nested sum involves factorial, exponentiation and greatest common divisor (gcd) computations. He further established that this formula also enumerates the number of unlabeled (n, r) -bipartite graphs, i.e., $|B_u(n, r)|$. A number of results indirectly related to Harrison's work and our result appeared in the literature [3, 4, 5, 6]. In particular, the set $B_u(n, r)$ in our work coincides with the set of bicolored graphs described in Section 2 in [3]. Whereas [3] provides a counting polynomial for the number of bicolored graphs, we focus on the asymptotic behavior of $|B_u(n, r)|$ in this paper. Counting polynomials for other families of bipartite graphs were also reported in [4]. Likewise, [5, 6] provide generating functions for related bipartite graph counting problems without an asymptotic analysis as provided in this paper. The species and category

*Department of Computer Engineering, Bilkent University, Ankara, Turkey.

†Department of Electrical and Computer Engineering, University of Maryland, College Park, MD 20742.

theory approach in [6] leads to a summation formula for the number of unlabeled bipartite graphs with v vertices. This formula is similar to the expression in (8) in [2] except that the latter formula counts the number of unlabeled bipartite graphs whose vertices are divided into two disjoint sets as in the model that is used in this paper. As such, for fixed n and r , the set $B_u(n, r)$ forms a subset of the set of unlabeled bipartite graphs with v vertices that are counted in [5, 6], where $v = n + r$. It should also be mentioned that some results on asymptotic enumeration of certain families of bipartite graphs (binary matrices) have been reported (see for example, [7, 8, 9, 10]). To the best of our knowledge, our work provides the first asymptotic enumeration of unlabelled bipartite graphs.

That $|B_u(1, r)| = r + 1$ trivially holds. Exact closed form expressions for $|B_u(n, r)|$ for $n = 2$, $n = 3$, and any integer $r > n$ are also given elsewhere[11]. The main result of this paper is the proof of the two-sided inequality given in (1).

Let S_n denote the symmetric group of permutations of degree n acting on set $N = \{1, 2, \dots, n\}$. Suppose that the $n!$ permutations in S_n are indexed by $1, 2, \dots, n!$ in some arbitrary, but fixed manner. The cycle index polynomial of S_n is defined as follows([12],see p.35, Eqn. 2.2.1):

$$Z_{S_n}(x_1, x_2, \dots, x_n) = \frac{1}{n!} \sum_{m=1}^{n!} \prod_{k=1}^n x_k^{p_{m,k}} \quad (2)$$

where $p_{m,k}$ denotes the number of cycles of length k in the disjoint cycle representation of the m^{th} permutation in S_n , and $\sum_{k=1}^n kp_{m,k} = n, \forall m = 1, 2, \dots, n!$.

Let $S_n \times S_r$ denote the direct product of symmetric groups S_n and S_r acting on $N = \{1, 2, \dots, n\}$ and $R = \{1, 2, \dots, r\}$, respectively, where n and r are positive integers such that $n < r$. It can be inferred from Harrison ([13],Lemma 4.1 and Theorem 4.2) that the cycle index polynomial of $S_n \times S_r$ is given by [13]

$$Z_{S_n \times S_r}(x_1, x_2, \dots, x_{nr}) = Z_{S_n}(x_1, x_2, \dots, x_n) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r), \quad (3)$$

where \boxtimes is a particular polynomial multiplication that distributes over ordinary addition, and in which the multiplication $X_m \odot X_t$ of two product terms¹, $X_m = x_1^{p_{m,1}} x_2^{p_{m,2}} \dots x_n^{p_{m,n}}$ and $X_t = x_1^{q_{t,1}} x_2^{q_{t,2}} \dots x_r^{q_{t,r}}$ in Z_{S_n} and Z_{S_r} , respectively, is defined as²

$$X_m \odot X_t = \prod_{k=1}^n \prod_{j=1}^r x_{\text{lcm}(k,j)}^{p_{m,k} q_{t,j} \text{gcd}(k,j)}. \quad (4)$$

Harrison further proved that [2]:

$$|B_u(n, r)| = Z_{S_n \times S_r}(\underbrace{2, 2, \dots, 2}_{nr}) \quad (5)$$

when³ $n \neq r$.

We need one more fact that can be found in Harary ([12], p.36) in order to compute the stated lower and upper bound in (1):

$$Z_{S_r}(x_1, x_2, \dots, x_r) = \frac{1}{r} \sum_{i=1}^r x_i Z_{S_{r-i}}(x_1, x_2, \dots, x_{r-i}) \quad (6)$$

where $Z_{S_0}() = 1$.

¹Note that we will not display the zero powers of x_1, x_2, \dots in a cycle index polynomial. We will use the same convention for all other cycle index polynomials throughout the paper.

²The $\text{lcm}(a,b)$ and $\text{gcd}(a,b)$ denote least common multiple and greatest common divisor of a and b .

³As noted in [2], $n = r$ case involves a different cycle index polynomial. Bounding $|B_u(n, n)|$ will be considered separately at the end of the paper.

2 The Lower Bound for $|B_u(n, r)|$

From (3) and (5) we know that

$$|B_u(n, r)| = Z_{S_n \times S_r}(2, 2, \dots, 2), \quad (7)$$

$$= [Z_{S_n}(x_1, x_2, \dots, x_n) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2). \quad (8)$$

One of the terms in $Z_{S_n}(x_1, x_2, \dots, x_n)$ is $\frac{1}{n!}(x_1^n)$ and it is associated with the identity permutation in S_n . Using this fact, we find

$$|B_u(n, r)| = Z_{S_n \times S_r}(2, 2, \dots, 2), \quad (9)$$

$$= [Z_{S_n}(x_1, x_2, \dots, x_n) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2), \quad (10)$$

$$= \left[\left(\frac{1}{n!}(x_1^n + \dots) \right) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r) \right] (2, 2, \dots, 2), \quad (11)$$

$$= \left[\left(\frac{1}{n!}x_1^n \right) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r) \right] (2, 2, \dots, 2) + \dots, \quad (12)$$

$$= \frac{1}{n!} \left\{ \left[x_1^n \boxtimes \frac{1}{r!} \sum_{t=1}^{r!} \prod_{j=1}^r x_j^{q_{t,j}} \right] (2, 2, \dots, 2) \right\} + \dots, \quad (13)$$

$$= \frac{1}{n!} \left\{ \left[\frac{1}{r!} \sum_{t=1}^{r!} x_1^n \odot \prod_{j=1}^r x_j^{q_{t,j}} \right] (2, 2, \dots, 2) \right\} + \dots, \quad (14)$$

$$= \frac{1}{n!} \left\{ \left[\frac{1}{r!} \sum_{t=1}^{r!} \prod_{j=1}^r x_j^{nq_{t,j} \gcd(1,j)} \right] (2, 2, \dots, 2) \right\} + \dots, \quad (15)$$

$$= \frac{1}{n!} \left\{ \left[\frac{1}{r!} \sum_{t=1}^{r!} \prod_{j=1}^r x_j^{nq_{t,j}} \right] (2, 2, \dots, 2) \right\} + \dots, \quad (16)$$

$$= \frac{1}{n!} \left\{ \frac{1}{r!} \sum_{t=1}^{r!} \prod_{j=1}^r 2^{nq_{t,j}} \right\} + \dots, \quad (17)$$

$$= \frac{1}{n!} \left\{ \frac{1}{r!} \sum_{t=1}^{r!} \prod_{j=1}^r (2^n)^{q_{t,j}} \right\} + \dots, \quad (18)$$

$$= \frac{1}{n!} \left\{ Z_{S_r}(2^n, 2^n, \dots, 2^n) \right\} + \dots \quad (19)$$

This proves

$$|B_u(n, r)| \geq \frac{1}{n!} Z_{S_r}(2^n, 2^n, \dots, 2^n). \quad \square \quad (20)$$

Proposition 1.

$$Z_{S_r}(2^n, 2^n, \dots, 2^n) = \binom{r + 2^n - 1}{r}$$

Proof. Using (6), we have

$$r Z_{S_r}(2^n, 2^n, \dots, 2^n) = \sum_{i=1}^r 2^n Z_{S_{r-i}}(2^n, 2^n, \dots, 2^n), \quad (21)$$

and

$$(r-1) Z_{S_{r-1}}(2^n, 2^n, \dots, 2^n) = \sum_{i=1}^{r-1} 2^n Z_{S_{r-1-i}}(2^n, 2^n, \dots, 2^n). \quad (22)$$

Subtracting the second equation from the first one gives

$$rZ_{S_r}(2^n, 2^n, \dots, 2^n) - (r-1)Z_{S_{r-1}}(2^n, 2^n, \dots, 2^n) = 2^n Z_{S_{r-1}}(2^n, 2^n, \dots, 2^n), \quad (23)$$

$$rZ_{S_r}(2^n, 2^n, \dots, 2^n) = (r+2^n-1)Z_{S_{r-1}}(2^n, 2^n, \dots, 2^n), \quad (24)$$

$$Z_{S_r}(2^n, 2^n, \dots, 2^n) = \left(\frac{r+2^n-1}{r}\right)Z_{S_{r-1}}(2^n, 2^n, \dots, 2^n). \quad (25)$$

Expanding the last equation inductively, we obtain

$$Z_{S_r}(2^n, 2^n, \dots, 2^n) = \left(\frac{r+2^n-1}{r}\right)\left(\frac{r+2^n-2}{r-1}\right)Z_{S_{r-2}}(2^n, 2^n, \dots, 2^n), \quad (26)$$

$$Z_{S_r}(2^n, 2^n, \dots, 2^n) = \left(\frac{r+2^n-1}{r}\right)\left(\frac{r+2^n-2}{r-1}\right)\left(\frac{r+2^n-3}{r-2}\right)Z_{S_{r-3}}(2^n, 2^n, \dots, 2^n), \quad (27)$$

$$Z_{S_r}(2^n, 2^n, \dots, 2^n) = \left(\frac{r+2^n-1}{r}\right)\left(\frac{r+2^n-2}{r-1}\right)\left(\frac{r+2^n-3}{r-2}\right)\dots\left(\frac{2^n}{1}\right)Z_{S_0}(). \quad (28)$$

Noting that $Z_{S_0}() = 1$, and combining the product terms together, we obtain

$$Z_{S_r}(2^n, 2^n, \dots, 2^n) = \binom{r+2^n-1}{r}. \quad \square \quad (29)$$

Combining Proposition 1 with (20) proves the lower bound:

Theorem 1.

$$|B_u(n, r)| \geq \frac{1}{n!}Z_{S_r}(2^n, 2^n, \dots, 2^n) \geq \frac{\binom{r+2^n-1}{r}}{n!}. \quad \square \quad (30)$$

3 An Upper Bound for $|B_u(n, r)|$

We first note that $|B_u(1, r)| = r+1 = \binom{r+2^1-1}{r}/1! \leq 2\binom{r+2^1-1}{r}/1!$. Hence the upper bound that is claimed in the abstract holds for $n=1$. Proving that it also holds for $n \geq 2$ requires a more careful analysis of the terms in

$$Z_{S_n}(x_1, x_2, \dots, x_n) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r). \quad (31)$$

We first express $Z_{S_n}(x_1, x_2, \dots, x_n)$ as

$$Z_{S_n}(x_1, x_2, \dots, x_n) = Z_{S_n}[1] + Z_{S_n}[2] + \dots + Z_{S_n}[n!], \quad (32)$$

where

$$Z_{S_n}[1] = \frac{1}{n!}x_1^n \quad (33)$$

$$Z_{S_n}[2] = \frac{1}{n!}x_1^{n-2}x_2 \quad (34)$$

The first term is associated with the identity permutation and the second term is associated with any one of the permutations in which all but two of the elements in $N = 1, 2, \dots, n$ are fixed to themselves. The remaining $Z_{S_n}[i] = \frac{1}{n!} \prod_{k=1}^n x_k^{p_{i,k}}, 3 \leq i \leq n!$ terms represent all the other product terms in the cycle index polynomial of S_n with no particular association with the permutations in S_n . Similarly, we set $Z_{S_r}(x_1, x_2, \dots, x_r) = \frac{1}{r!} \sum_{t=1}^{r!} \prod_{j=1}^r x_j^{q_{t,j}}$ without identifying the actual product terms with any particular permutation in S_r .

The following equations obviously hold as the sum of the lengths of all the cycles in any cycle disjoint representation of a permutation in S_n and S_r must be n and r , respectively.

$$\sum_{k=1}^n kp_{i,k} = n, 1 \leq i \leq n!, \quad (35)$$

$$\sum_{j=1}^r jq_{t,j} = r, 1 \leq t \leq r! \quad (36)$$

Now we can proceed with the computation of the upper bound for $|B_u(n, r)|$. First, we note that

$$|B_u(n, r)| = Z_{S_n \times S_r}(2, 2, 2, \dots, 2), \quad (37)$$

$$= [Z_{S_n}(x_1, x_2, \dots, x_n) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2), \quad (38)$$

$$= [(Z_{S_n}[1] + Z_{S_n}[2] + \dots + Z_{S_n}[n!]) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2), \quad (39)$$

$$= [Z_{S_n}[1] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) + [Z_{S_n}[2] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) \\ + \dots + [Z_{S_n}[n!] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2). \quad (40)$$

The first term in (40) is directly computed from Proposition 1. Thus, it suffices to upper bound each of the remaining terms in (40) to upper bound $|B_u(n, r)|$. This will be established by proving $[Z_{S_n}[2] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) \geq [Z_{S_n}[i] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2), \forall i, 3 \leq i \leq n!$. We first need some preliminary facts.

Lemma 1. For all $i, 1 \leq i \leq n!$,

$$[Z_{S_n}[i] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, \dots, 2) = \frac{1}{n!} Z_{S_r}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, \dots, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)}). \quad (41)$$

Proof.

$$[Z_{S_n}[i] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) = \left[\frac{1}{n!} \prod_{k=1}^n x_k^{p_{i,k}} \boxtimes \left(\frac{1}{r!} \sum_{t=1}^{r!} \prod_{j=1}^r x_j^{q_{t,j}} \right) \right] (2, 2, \dots, 2), \quad (42)$$

$$= \left[\frac{1}{n! r!} \sum_{t=1}^{r!} \prod_{k=1}^n x_k^{p_{i,k}} \odot \prod_{j=1}^r x_j^{q_{t,j}} \right] (2, 2, \dots, 2), \quad (43)$$

$$= \left[\frac{1}{n! r!} \sum_{t=1}^{r!} \prod_{j=1}^r \prod_{k=1}^n x_{\text{lcm}(k,j)}^{p_{i,k} q_{t,j} \gcd(k,j)} \right] (2, 2, \dots, 2), \quad (44)$$

$$= \frac{1}{n! r!} \sum_{t=1}^{r!} \prod_{j=1}^r \prod_{k=1}^n 2^{p_{i,k} q_{t,j} \gcd(k,j)}, \quad (45)$$

$$= \frac{1}{n!} \left[\frac{1}{r!} \sum_{t=1}^{r!} \prod_{j=1}^r (2^{\sum_{k=1}^n p_{i,k} \gcd(k,j)})^{q_{t,j}} \right], \quad (46)$$

$$= \frac{1}{n!} Z_{S_r}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, \dots, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)}). \quad \square \quad (47)$$

Corollary 1.

$$[Z_{S_n}[2] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, \dots, 2) = \frac{1}{n!} Z_{S_r}(2^{n-1}, 2^n, 2^{n-1}, 2^n, \dots). \quad (48)$$

Proof. By definition, $p_{2,1} = n - 2, p_{2,2} = 1, p_{2,k} = 0, 3 \leq k \leq n$. Substituting these into the last equation in Lemma 1 proves the statement. \square

Lemma 2.

$$\sum_{k=1}^n p_{i,k} \leq n - 1, \forall i, 2 \leq i \leq n!. \quad (49)$$

Proof. Recall from (35) that $\sum_{k=1}^n k p_{i,k} = n, \forall i, 1 \leq i \leq n!$. Hence $\sum_{k=1}^n p_{i,k} = n - \sum_{k=1}^n (k-1) p_{i,k}$, and so the maximum value of $\sum_{k=1}^n p_{i,k}$ occurs when $\sum_{k=1}^n (k-1) p_{i,k}$ is minimized. Furthermore, at least one of $p_{i,k}, \forall i, 2 \leq i \leq n!$ must be ≥ 1 for some $k \geq 2$ since none of the permutations we consider is the identity. Thus, $\sum_{k=1}^n (k-1) p_{i,k} \geq 1$ and the statement follows. \square

Lemma 3. If $\sum_{k=1}^n p_{i,k} \gcd(k, \alpha + 1) = n$, then $\sum_{k=1}^n p_{i,k} \gcd(k, \alpha) \leq n - 1, \forall i, 2 \leq i \leq n!$ and for any integer $\alpha \geq 2$.

Proof. If $\sum_{k=1}^n p_{i,k} \gcd(k, \alpha + 1) = n$ as stated in the lemma, then we must have $\gcd(k, \alpha + 1) = k$ where $p_{i,k} \geq 1, \forall i, 2 \leq i \leq n!$. Therefore $k \leq \alpha + 1$. Now if $k = \alpha + 1$, then trivially $\gcd(k, \alpha) < k$. On the other hand if $k < \alpha + 1$, then $\alpha + 1$ must be a multiple of k . Therefore, α can not be a multiple

of k for any $k \geq 2$. At this point we find that $\gcd(k, \alpha) < k, \forall k, 2 \leq k \leq n$. Since as in the previous lemma, none of the permutations we consider is the identity, at least one of $p_{i,k}, \forall i, 2 \leq i \leq n!$ must be ≥ 1 for some $k \geq 2$ and so we conclude that $\sum_{k=1}^n p_{i,k} \gcd(k, \alpha) \leq n - 1$. \square

Lemma 4. $Z_{S_r}(2^{n-1}, 2^n, \dots) \geq Z_{S_{r-1}}(2^{n-1}, 2^n, \dots)$, for $2 \leq n$.

Proof. Using (6), we get

$$rZ_{S_r}(2^{n-1}, 2^n, \dots) = \sum_{\text{odd } i}^{r-\beta_1} 2^{n-1} Z_{S_{r-i}}(2^{n-1}, 2^n, \dots) + \sum_{\text{even } i}^{r-\beta_2} 2^n Z_{S_{r-i}}(2^{n-1}, 2^n, \dots), \quad (50)$$

where $\beta_1 = 1, \beta_2 = 0$ if r is even and $\beta_1 = 0, \beta_2 = 1$ if r is odd. Similarly, for $r - 1$,

$$(r-1)Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) = \sum_{\text{odd } i}^{r-1-\beta_2} 2^{n-1} Z_{S_{r-1-i}}(2^{n-1}, 2^n, \dots) + \sum_{\text{even } i}^{r-1-\beta_1} 2^n Z_{S_{r-1-i}}(2^{n-1}, 2^n, \dots). \quad (51)$$

Subtracting 51 from 50 gives

$$\begin{aligned} & rZ_{S_r}(2^{n-1}, 2^n, \dots) - (r-1)Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) \\ &= \sum_{\text{even } i}^{r-\beta_2} 2^n Z_{S_{r-i}}(2^{n-1}, 2^n, \dots) - \sum_{\text{odd } i}^{r-1-\beta_2} 2^{n-1} Z_{S_{r-1-i}}(2^{n-1}, 2^n, \dots) \\ & \quad + \sum_{\text{odd } i}^{r-\beta_1} 2^{n-1} Z_{S_{r-i}}(2^{n-1}, 2^n, \dots) - \sum_{\text{even } i}^{r-1-\beta_1} 2^n Z_{S_{r-1-i}}(2^{n-1}, 2^n, \dots), \end{aligned} \quad (52)$$

$$\begin{aligned} & rZ_{S_r}(2^{n-1}, 2^n, \dots) - (r-1)Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) \\ &= \sum_{\text{even } i}^{r-\beta_2} 2^{n-1} Z_{S_{r-i}}(2^{n-1}, 2^n, \dots) + 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) - \sum_{\text{even } i}^{r-1-\beta_1} 2^{n-1} Z_{S_{r-1-i}}(2^{n-1}, 2^n, \dots), \end{aligned} \quad (53)$$

$$\begin{aligned} & rZ_{S_r}(2^{n-1}, 2^n, \dots) = (r-1 + 2^{n-1})Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) \\ & \quad + 2^{n-1} \left(\sum_{\text{even } i}^{r-\beta_2} Z_{S_{r-i}}(2^{n-1}, 2^n, \dots) - \sum_{\text{even } i}^{r-1-\beta_1} Z_{S_{r-1-i}}(2^{n-1}, 2^n, \dots) \right). \end{aligned} \quad (54)$$

We now prove the lemma by induction on r .

Basis $r = 1$. By (6), $Z_{S_1}(2^{n-1}) = 2^{n-1} Z_{S_0}() = 2^{n-1}$. So we have $Z_{S_1}(2^{n-1}) = 2^{n-1} \geq Z_{S_0}() = 1$ for $2 \leq n$.

Induction Step. Suppose that the lemma holds from 1 to $r - 1$. That is, $Z_{S_{r-i}} - Z_{S_{r-i-1}} \geq 0, 1 \leq i \leq r - 1$. Now if r is even then the difference of the two sums in (54) becomes $(Z_{S_{r-2}} - Z_{S_{r-3}}) + (Z_{S_{r-4}} - Z_{S_{r-5}}) \dots + (Z_{S_2} - Z_{S_1}) + Z_{S_0}$, which is clearly ≥ 0 by the induction hypothesis. Therefore,

$$rZ_{S_r}(2^{n-1}, 2^n, \dots) \geq (r-1 + 2^{n-1})Z_{S_{r-1}}(2^{n-1}, 2^n, \dots), \quad (55)$$

$$Z_{S_r}(2^{n-1}, 2^n, \dots) \geq Z_{S_{r-1}}(2^{n-1}, 2^n, \dots), n \geq 2. \quad (56)$$

On the other hand, if r is odd then the difference of the two sums in the same equation becomes $(Z_{S_{r-2}} - Z_{S_{r-3}}) + (Z_{S_{r-4}} - Z_{S_{r-5}}) \dots + (Z_{S_2} - Z_{S_1}) + (Z_{S_1} - Z_{S_0})$, which is again ≥ 0 , and the statement follows in this case as well. \square

We now are ready to prove that

$$[Z_{S_n}[2] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, \dots, 2) \geq [Z_{S_n}[i] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, \dots, 2), \forall i, 2 \leq i \leq n!.$$

Theorem 2.

$$[Z_{S_n}[2] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) \geq [Z_{S_n}[i] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) \quad (57)$$

$\forall i, 2 \leq i \leq n!$ and $\forall n, n < r$.

Proof. Using Lemma 1 and Corollary 1 it suffices to show that

$$Z_{S_r}(2^{n-1}, 2^n, \dots) \geq Z_{S_r}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, \dots, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)}). \quad (58)$$

We prove the statement by induction on r .

Basis: ($r = 1$). By (6), $Z_{S_1}(2^{n-1}) = 2^{n-1} Z_{S_0}() = 2^{n-1}$. Similarly, by (6), $Z_{S_1}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}) = 2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)} Z_{S_0}() = 2^{\sum_{k=1}^n p_{i,k}}$. Given that $\sum_{k=1}^n p_{i,k} \leq n - 1$ by Lemma 2, we have $2^{\sum_{k=1}^n p_{i,k}} \leq 2^{n-1}$, and hence the statement holds in this case.

Induction Step: First, by (6),

$$Z_{S_r}(2^{n-1}, 2^n, \dots) = \frac{1}{r} \begin{bmatrix} 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) \\ + 2^n Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) \\ + 2^{n-1} Z_{S_{r-3}}(2^{n-1}, 2^n, \dots) \\ \vdots \\ + 2^\beta Z_{S_0}() \end{bmatrix}, \quad (59)$$

where $\beta = n$ if r is even and $\beta = n - 1$ if r is odd. Similarly,

$$Z_{S_r}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, \dots, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)}) = \frac{1}{r} \begin{bmatrix} 2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)} Z_{S_{r-1}}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, \dots) \\ + 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)} Z_{S_{r-2}}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, \dots) \\ + 2^{\sum_{k=1}^n p_{i,k} \gcd(k,3)} Z_{S_{r-3}}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, \dots) \\ \vdots \\ + 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)} Z_{S_0}() \end{bmatrix} \quad (60)$$

Subtracting (60) from (59), we have

$$\begin{aligned} & Z_{S_r}(2^{n-1}, 2^n, \dots) - Z_{S_r}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, \dots, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)}) \\ &= \frac{1}{r} \begin{bmatrix} 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) \\ + 2^n Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) \\ + 2^{n-1} Z_{S_{r-3}}(2^{n-1}, 2^n, \dots) \\ \vdots \\ + 2^\beta Z_{S_0}() \end{bmatrix} - \frac{1}{r} \begin{bmatrix} 2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)} Z_{S_{r-1}}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)}, \dots) \\ + 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)} Z_{S_{r-2}}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)}, \dots) \\ + 2^{\sum_{k=1}^n p_{i,k} \gcd(k,3)} Z_{S_{r-3}}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)}, \dots) \\ \vdots \\ + 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)} Z_{S_0}() \end{bmatrix} \end{aligned} \quad (61)$$

Thus, it suffices to show that the right hand side of the above equation is ≥ 0 , or

$$\begin{aligned} & 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)} Z_{S_{r-1}}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)}, \dots) \\ & + 2^n Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)} Z_{S_{r-2}}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)}, \dots) \\ & + 2^{n-1} Z_{S_{r-3}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,3)} Z_{S_{r-3}}(2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)}, 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)}, \dots) \\ & \quad \vdots \\ & + 2^\beta Z_{S_0}() - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)} Z_{S_0}() \geq 0. \end{aligned} \quad (62)$$

Now by induction hypothesis, (58) holds for $1, 2, \dots, r - 1$. Thus, (62) can be replaced by

$$\begin{aligned}
& 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) \\
& + 2^n Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)} Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) \\
& + 2^{n-1} Z_{S_{r-3}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,3)} Z_{S_{r-3}}(2^{n-1}, 2^n, \dots) \\
& \quad \vdots \\
& + 2^\beta Z_{S_0}() - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)} Z_{S_0}() \geq 0.
\end{aligned} \tag{63}$$

Moreover, invoking Lemma 2 gives

$$\begin{aligned}
& 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,1)} Z_{S_{r-1}}(2^{n-1}, 2^n \dots) \\
& \geq 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) - 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) = 0.
\end{aligned} \tag{64}$$

Hence the difference in the first line in (63) ≥ 0 , and therefore it is sufficient to show that

$$\begin{aligned}
& 2^n Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,2)} Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) \\
& + 2^{n-1} Z_{S_{r-3}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,3)} Z_{S_{r-3}}(2^{n-1}, 2^n, \dots) \\
& \quad \vdots \\
& + 2^\beta Z_{S_0}() - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)} Z_{S_0}() \geq 0.
\end{aligned} \tag{65}$$

To prove this inequality, we will combine four terms in pairs of consecutive lines for the remaining $r - 1$ lines by considering two cases. If r is odd then $\beta = n - 1$ and no extra line remains in this pairing. Thus, for all even $\alpha, 2 \leq \alpha \leq r - 1$, it suffices to prove

$$\begin{aligned}
& 2^n Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,\alpha)} Z_{S_{r-\alpha}}(2^{n-1}, 2^n \dots), \\
& + 2^{n-1} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,\alpha+1)} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n \dots) \geq 0.
\end{aligned} \tag{66}$$

or,

$$\begin{aligned}
& 2^n Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} k} Z_{S_{r-\alpha}}(2^{n-1}, 2^n \dots) \\
& + 2^{n-1} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,\alpha+1)} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n \dots) \geq 0.
\end{aligned} \tag{67}$$

Now if $\sum_{k=1}^n p_{i,k} \gcd(k, \alpha + 1) \leq n - 1$, then

$$\begin{aligned}
& 2^n Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) - 2^{\sum_{k=1}^n p_{i,k} k} Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) \\
& + 2^{n-1} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots) - 2^{n-1} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots) \geq \\
& 2^n Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) - 2^n Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) \\
& + 2^{n-1} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots) - 2^{n-1} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots) = 0.
\end{aligned} \tag{68}$$

On the other hand, if $\sum_{k=1}^n p_{i,k} \gcd(k, \alpha + 1) = n$, then we prove (66) by noting that $\sum_{k=1}^n p_{i,k} \gcd(k, \alpha) \leq n - 1$ by Lemma 3. Thus,

$$2^n Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) - 2^{n-1} Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) \tag{69}$$

$$\begin{aligned}
& + 2^{n-1} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots) - 2^n Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots) \\
& = 2^{n-1} Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) - 2^{n-1} Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots) \\
& 2^{n-1} [Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) - Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots)]
\end{aligned} \tag{70}$$

Now by Lemma 4, $Z_{S_{r-\alpha}}(2^{n-1}, 2^n, \dots) \geq Z_{S_{r-\alpha-1}}(2^{n-1}, 2^n, \dots)$ and the statement is proved for odd $r, n < r$. For even r , the last line in (65) is left out in the pairing of consecutive lines and $\beta = n$. In this case we have $2^n Z_{S_0}() - 2^{\sum_{k=1}^n p_{i,k} \gcd(k,r)} Z_{S_0}() \geq 2^n Z_{S_0}() - 2^{\sum_{k=1}^n p_{i,k} k} Z_{S_0}() = 2^n Z_{S_0}() - 2^n Z_{S_0}() = 0$ and the statement follows. \square

Theorem 3.

$$[Z_{S_n}[2] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) \leq \frac{\binom{r+2^n-1}{r}}{n!(n!-1)}. \quad (71)$$

where $2 \leq n < r$.

Proof. By Corollary 1

$$[Z_{S_n}[2] \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) = \frac{1}{n!} Z_{S_r}(2^{n-1}, 2^n, \dots). \quad (72)$$

Thus, to prove the theorem, it is sufficient to show

$$\frac{1}{n!} Z_{S_r}(2^{n-1}, 2^n, 2^{n-1}, 2^n, \dots) \leq \frac{\binom{r+2^n-1}{r}}{n!(n!-1)} \quad (73)$$

where $2 \leq n < r$.

Now, using (6), we get

$$r Z_{S_r}(2^{n-1}, 2^n, \dots) = \sum_{\text{odd } i}^{r-\beta_1} 2^{n-1} Z_{S_{r-i}}(2^{n-1}, 2^n, \dots) + \sum_{\text{even } i}^{r-\beta_2} 2^n Z_{S_{r-i}}(2^{n-1}, 2^n, \dots) \quad (74)$$

where $\beta_1 = 1, \beta_2 = 0$ if r is even and $\beta_1 = 0, \beta_2 = 1$ if r is odd. Similarly, for $r-2$,

$$(r-2) Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) = \sum_{\text{odd } i}^{r-2-\beta_1} 2^{n-1} Z_{S_{r-2-i}}(2^{n-1}, 2^n, \dots) + \sum_{\text{even } i}^{r-2-\beta_2} 2^n Z_{S_{r-2-i}}(2^{n-1}, 2^n, \dots). \quad (75)$$

Subtracting (75) from (74) gives

$$\begin{aligned} & r Z_{S_r}(2^{n-1}, 2^n, \dots) - (r-2) Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) \\ &= 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) + 2^n Z_{S_{r-2}}(2^{n-1}, 2^n, \dots), \end{aligned} \quad (76)$$

$$r Z_{S_r}(2^{n-1}, 2^n, \dots) = 2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) + (r-2+2^n) Z_{S_{r-2}}(2^{n-1}, 2^n, \dots), \quad (77)$$

$$Z_{S_r}(2^{n-1}, 2^n, \dots) = \frac{1}{r} [2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) + (r-2+2^n) Z_{S_{r-2}}(2^{n-1}, 2^n, \dots)]. \quad (78)$$

We will use induction on r and the recurrence given in (78) to prove this inequality.

Basis. Case $r = 3$: Recall that

$$Z_{S_n}[2] = \frac{1}{n!} x_1^{n-2} x_2, \quad (79)$$

$$Z_{S_3}(x_1, x_2, x_3) = \frac{1}{3!} (x_1^3 + 3x_1 x_2 + 2x_3). \quad (80)$$

Thus,

$$\begin{aligned} & [Z_{S_n}[2] \boxtimes Z_{S_3}(x_1, x_2, x_3)](2, 2, \dots, 2) \\ &= \left[\frac{1}{n!} (x_1^{n-2} x_2) \boxtimes \frac{1}{3!} (x_1^3 + 3x_1 x_2 + 2x_3) \right] (2, 2, \dots, 2), \end{aligned} \quad (81)$$

$$= \frac{1}{3!n!} \left[(x_1^{n-2} x_2) \odot x_1^3 + (x_1^{n-2} x_2) \odot (3x_1 x_2) + (x_1^{n-2} x_2) \odot 2x_3 \right] (2, 2, \dots, 2), \quad (82)$$

$$= \frac{1}{3!n!} \left[x_1^{3(n-2)} x_2^3 + 3x_1^{n-2} x_2 x_2^{n-2} x_2^2 + 2x_3^{n-2} x_6 \right] (2, 2, \dots, 2), \quad (83)$$

$$= \frac{1}{3!n!} [2^{3n-3} + 3 \times 2^{2n-1} + 2^n] \leq \frac{\binom{r+2^n-1}{r}}{n!(n!-1)}. \quad (84)$$

for $n = 2$ and $r = 3$.

Case $r = 4$. In this case we have

$$\begin{aligned} & [Z_{S_n}[2] \boxtimes Z_{S_4}(x_1, x_2, x_3, x_4)](2, \dots, 2) \\ &= \left[\frac{1}{n!} (x_1^{n-2} x_2) \boxtimes \frac{1}{4!} (x_1^4 + 6x_1^2 x_2 + 3x_2^2 + 8x_1 x_3 + 6x_4) \right] (2, \dots, 2), \end{aligned} \quad (85)$$

$$\begin{aligned} &= \frac{1}{4!n!} \left[(x_1^{n-2} x_2) \odot x_1^4 + (x_1^{n-2} x_2) \odot (6x_1^2 x_2) + (x_1^{n-2} x_2) \odot 3x_2^2 \right. \\ &\quad \left. + (x_1^{n-2} x_2) \odot (8x_1 x_3) + (x_1^{n-2} x_2) \odot 6x_4 \right] (2, \dots, 2), \end{aligned} \quad (86)$$

$$= \frac{1}{4!n!} \left[x_1^{4(n-2)} x_2^4 + 6x_1^{2(n-2)} x_2^2 x_2^2 + 3x_1^{2(n-2)} x_2^4 + 8x_1^{n-2} x_3^{n-2} x_2 x_6 + 6x_4^{n-2} x_4^2 \right] (2, \dots, 2), \quad (87)$$

$$= \frac{1}{4!n!} \left[2^{4n-4} + 6 \times 2^{3n-2} + 3 \times 2^{2n} + 8 \times 2^{2n-2} + 6 \times 2^n \right], \quad (88)$$

$$= \frac{1}{4!n!} \left[2^{4n-4} + 6 \times 2^{3n-2} + 5 \times 2^{2n} + 6 \times 2^n \right]. \quad (89)$$

Now, given that $r = 4$, the only possible values of n are 2 and 3. If $n = 2$ then:

$$[Z_{S_n}[2] \boxtimes Z_{S_4}(x_1, x_2, x_3, x_4)](2, 2, \dots, 2) = \frac{1}{4!n!} \left[2^{4n-4} + 6 \times 2^{3n-2} + 5 \times 2^{2n} + 6 \times 2^n \right], \quad (90)$$

$$= \frac{1}{4!2!} \left[2^4 + 6 \times 2^4 + 5 \times 2^4 + 6 \times 2^2 \right], \quad (91)$$

$$= \frac{16 + 96 + 80 + 24}{4!2!} = 4.5, \quad (92)$$

$$\leq \frac{\binom{r+2^n-1}{r}}{n!(n!-1)} = \frac{\binom{7}{4}}{2!(2!-1)} = \frac{35}{2} = 17.5. \quad (93)$$

On the other hand, if $n = 3$ then:

$$[Z_{S_n}[2] \boxtimes Z_{S_4}(x_1, x_2, x_3, x_4)](2, 2, \dots, 2) = \frac{1}{4!n!} \left[2^{4n-4} + 6 \times 2^{3n-2} + 5 \times 2^{2n} + 6 \times 2^n \right], \quad (94)$$

$$= \frac{1}{4!3!} \left[2^8 + 6 \times 2^7 + 5 \times 2^6 + 6 \times 2^3 \right], \quad (95)$$

$$= \frac{256 + 768 + 320 + 48}{4!3!} = \frac{29}{3}, \quad (96)$$

$$\leq \frac{\binom{r+2^n-1}{r}}{n!(n!-1)} = \frac{\binom{11}{4}}{3!(3!-1)} = \frac{330}{30} = 11. \quad (97)$$

Induction Step: Suppose that (73) holds for all values from 3 to $r-1$. Using the recurrence given in (78) and the induction hypothesis for $r-1$ and $r-2$ we get:

$$\frac{1}{n!} Z_{S_r}(2^{n-1}, 2^n, \dots) = \frac{1}{n!r} \left[2^{n-1} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) + (r-2+2^n) Z_{S_{r-2}}(2^{n-1}, 2^n, \dots) \right], \quad (98)$$

$$= \frac{2^{n-1}}{n!r} Z_{S_{r-1}}(2^{n-1}, 2^n, \dots) + \frac{r-2+2^n}{n!r} Z_{S_{r-2}}(2^{n-1}, 2^n, \dots), \quad (99)$$

$$\leq \frac{2^{n-1}}{r} \frac{\binom{r+2^n-2}{r-1}}{n!(n!-1)} + \frac{r-2+2^n}{r} \frac{\binom{r+2^n-3}{r-2}}{n!(n!-1)}, \quad (100)$$

$$\leq \frac{2^{n-1}}{n!(n-1)r} \frac{(r+2^n-2)!}{(r-1)!(2^n-1)!} + \frac{r-2+2^n}{n!(n-1)r} \frac{(r+2^n-3)!}{(r-2)!(2^n-1)!}, \quad (101)$$

$$\leq \frac{2^{n-1}}{n!(n-1)r} \frac{(r+2^n-2)!}{(r-1)!(2^n-1)!} + \frac{(r-1)(r+2^n-2)!}{n!(n-1)r!(2^n-1)!}, \quad (102)$$

$$\frac{1}{n!} Z_{S_r}(2^{n-1}, 2^n, \dots) \leq \frac{(r+2^n-2)!(r+2^{n-1}-1)}{n!(n-1)r!(2^n-1)!} \leq \frac{(r+2^n-2)!(r+2^n-1)}{n!(n-1)r!(2^n-1)!}, \quad (103)$$

$$\leq \frac{(r+2^n-1)!}{n!(n-1)r!(2^n-1)!} = \frac{1}{n!(n-1)} \binom{r+2^n-1}{r}, \quad (104)$$

$$\leq \frac{1}{n!(n-1)} \binom{r+2^n-1}{r}. \quad (105)$$

This completes the proof. \square

Combining Theorems 2 and 3 concludes the upper bound calculation.

Theorem 4. $|B_u(n, r)| \leq \frac{2^{\binom{r+2^n-1}{r}}}{n!}$.

Proof.

$$|B_u(n, r)| = Z_{S_n \times S_r}(2, 2, \dots, 2), \quad (106)$$

$$= [Z_{S_n}(x_1, x_2, \dots, x_n) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2), \quad (107)$$

$$= [(Z_{S_n}[1] + Z_{S_n}[2] + \dots + Z_{S_n}[n!]) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2), \quad (108)$$

$$= [(Z_{S_n}[1]) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) + [(Z_{S_n}[2]) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) + \dots + [(Z_{S_n}[n!]) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2), \quad (109)$$

$$\leq [(Z_{S_n}[1]) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) + [(Z_{S_n}[2]) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2) + \dots + [(Z_{S_n}[n!]) \boxtimes Z_{S_r}(x_1, x_2, \dots, x_r)](2, 2, \dots, 2), \quad (110)$$

$$\leq \frac{\binom{r+2^n-1}{r}}{n!} + (n-1) \frac{\binom{r+2^n-1}{r}}{n!(n-1)} = \frac{2^{\binom{r+2^n-1}{r}}}{n!}. \quad \square \quad (111)$$

Remark 1. It should be mentioned that, if $r < n$, using the relation $|B_u(n, r)| = |B_u(r, n)|$ gives

$$|B_u(n, r)| \leq 2^{\frac{\binom{n+2^r-1}{n}}{r!}}. \quad (112)$$

Likewise, if $r < n$, Theorem 1 and $|B_u(n, r)| = |B_u(r, n)|$ together imply

$$|B_u(n, r)| \geq \frac{\binom{n+2^r-1}{n}}{r!}. \quad (113)$$

Furthermore, if $r = n$, using the cycle index representation of bi-colored graphs provided in Section 3 in [3] and Theorem 1 gives

$$|B_u(n, n)| \geq \frac{\binom{n+2^n-1}{n}}{2n!}. \quad (114)$$

The Z' term in the cycle index representation of bi-colored graphs in [3] prevents us from deriving an upper bound for $|B_u(n, n)|$ that is a constant multiple of the lower bound in this case. On the other hand, an obvious upper bound for $|B_u(n, n)|$ can be derived by setting $r = n + 1$ in the inequality in Theorem 4.

Appendix:

Table 1 lists $\ln |B_u(n, r)|$ along with the natural logarithms of lower and upper bounds for $1 \leq n < r \leq 15$.

n	r	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1		1.09861 1.09861 1.79176	1.38629 1.38629 2.07944	1.80944 1.80944 2.30259	1.79176 1.79176 2.48491	1.94591 1.94591 2.63906	2.07944 2.07944 2.77259	2.19722 2.19722 2.89037	2.30259 2.30259 2.99573	2.3979 2.3979 3.09104	2.48491 2.48491 3.17805	2.58495 2.58495 3.2581	2.63906 2.63906 3.3322	2.70805 2.70805 3.4012	2.77259 2.77259 3.46574
2			2.30259 2.56495 2.99573	2.83321 3.09104 3.55355	3.3322 3.52636 4.02535	3.73767 3.91202 4.43082	4.09434 4.2485 4.78749	4.40672 4.55388 5.10595	4.70048 4.82831 5.39363	4.96284 5.0614 5.65599	5.20401 5.31321 5.89715	5.42495 5.52943 6.1203	5.63479 5.7301 6.32794	5.82895 5.91889 6.52209	6.01127 6.09582 6.70441
3				4.00733 4.46591 4.70048	4.8828 5.24702 5.57595	5.65599 5.95584 6.34914	6.34914 6.59851 7.04229	6.97728 7.18841 7.67089	7.55276 7.73368 8.24617	8.08364 8.24012 8.77678	8.57622 8.71276 9.26936	9.03575 9.1562 9.7289	9.46653 9.57345 10.1597	9.872 9.96754 10.5651	10.255 10.3409 10.9481
4					6.4708 6.9594 7.16395	7.72356 8.08641 8.41671	8.86869 9.14238 9.56184	9.92471 10.1349 10.6179	10.9056 11.0692 11.5987	11.8219 11.9512 12.515	12.6821 12.7855 13.3752	13.493 13.5767 14.1861	14.2603 14.3287 14.9534	14.9885 15.045 15.6816	15.6816 15.7287 16.3748
5						9.87164 10.2603 10.5648	11.5633 13.3276 12.2565	13.1474 14.7645 13.8406	14.6391 16.1388 15.3322	16.0501 17.4535 16.7432	17.3899 18.7124 18.083	18.6662 19.9195 19.3593	19.8854 21.0784 20.5785	21.053 21.7461 21.7461	22.1736 22.1927 22.8667
6							14.3253 14.5771 15.0185	16.5086 16.6637 17.2017	18.588 18.6849 19.2811	20.5759 20.6372 21.269	22.482 22.5215 23.1752	24.3146 24.3403 25.0078	26.0804 26.0974 26.7736	27.7852 27.7965 28.4783	29.4338 29.4415 30.127
7								19.9011 20.0463 20.5942	22.6165 22.6996 23.3097	25.2339 25.282 25.927	27.7633 27.7915 28.4564	30.2128 30.2295 30.906	32.5895 32.5995 33.2827	34.8992 34.9053 35.5924	37.147 37.1507 37.8401
8								26.6393 26.7201 27.3324	29.9164 29.9604 30.6096	33.102 33.1261 33.7952	36.2043 36.2177 36.8975	39.2304 39.2378 39.9235	42.186 42.1902 42.8792	45.0764 45.0788 45.7696	
9								34.5644 34.6096 35.2575	38.4241 38.4479 39.1173	42.1988 42.2114 42.892	45.8953 45.902 46.5885	49.5197 49.5233 50.2128	53.0769 53.0789 53.7701		
10									43.693 43.7187 44.3861	48.1502 48.1635 48.8434	52.5284 52.5353 53.2216	56.8335 56.837 57.5266	61.0705 61.0723 61.7636		
11									54.0381 54.0528 54.7312	59.1036 59.1111 59.7967	64.0955 64.0993 64.7886	69.0189 69.0208 69.712			
12											65.6106 65.6191 66.3038	71.2925 71.2968 71.9856	76.9056 76.9078 77.5988		
13													78.4205 78.4254 79.1137	84.7251 84.7275 85.4182	
14														92.4768 92.4797 93.17	

Table 1: Exact values of $\ln |B_u(n, r)|$, $1 \leq n < r \leq 15$, and natural logarithms of lower and upper bounds.

References

- [1] A. Yavuz Oruc. *Foundations of interconnection networks. To be published.* CRC press, 2016.
- [2] Michael A Harrison. On the number of classes of binary matrices. *IEEE Transactions on Computers*, 22(12):1048–1052, 1973.
- [3] Frank Harary. On the number of bi-colored graphs. *Pacific J. Math*, 8(1958):743–755, 1958.
- [4] Frank Harary and Geert Prins. Enumeration of bicolourable graphs. *Canad. J. Math*, 15:237–248, 1963.
- [5] Phil Hanlon. The enumeration of bipartite graphs. *Discrete Mathematics*, 28(1):49–57, 1979.
- [6] Andrew Gainer-Dewar and Ira M Gessel. Enumeration of bipartite graphs and bipartite blocks. *The Electronic Journal of Combinatorics*, 21(2):P2–40, 2014.
- [7] E Rodney Canfield, Brendan D McKay, et al. Asymptotic enumeration of dense 0-1 matrices with equal row sums and equal column sums. *Journal of Combinatorics*, 12(2):R29, 2005.
- [8] Catherine Greenhill, Brendan D McKay, and Xiaoji Wang. Asymptotic enumeration of sparse 0–1 matrices with irregular row and column sums. *Journal of Combinatorial Theory, Series A*, 113(2):291–324, 2006.
- [9] E Rodney Canfield, Catherine Greenhill, and Brendan D McKay. Asymptotic enumeration of dense 0–1 matrices with specified line sums. *Journal of Combinatorial Theory, Series A*, 115(1):32–66, 2008.
- [10] Alexander Barvinok. On the number of matrices and a random matrix with prescribed row and column sums and 0–1 entries. *Advances in Mathematics*, 224(1):316–339, 2010.
- [11] A. Atmaca and A Yavuz Oruc. Exact closed-form formulas for the number of certain unlabeled bipartite graphs. To be published, 2016.
- [12] Frank Harary. *Graphical Enumeration.* ACADEMIC PRESS, INC, 1973.
- [13] Michael A Harrison. Counting theorems and their applications to classification of switching functions. *Recent developments in switching theory*, pages 85–120, 1971.